

Product datasheet for AR51049PU-N

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

MPZL1 (38-162, His-tag) Human Protein

Product data:

Product Type: Recombinant Proteins

Description: MPZL1 (38-162, His-tag) human protein, 0.25 mg

Species: Human
Expression Host: E. coli

Expression cDNA Clone MGSSHHHHHH SSGLVPRGSH MGSLEVYTPK EIFVANGTQG KLTCKFKSTS TTGGLTSVSW

or AA Sequence: SFQPEGADTT VSFFHYSQGQ VYLGNYPPFK DRISWAGDLD KKDASINIEN MQFIHNGTYI

CDVKNPPDIV VQPGHIRLYV VEKENLPV

Tag: His-tag

Predicted MW: 16.1 kDa

Concentration: lot specific

Purity: >90% by SDS - PAGE

Buffer: Presentation State: Purified

State: Liquid purified protein

Buffer System: 20 mM Tris-HCl buffer (pH 8.0) containing 0.15M NaCl, 20% glycerol, 1 mM

DTT

Preparation: Liquid purified protein

Storage: Store undiluted at 2-8°C for one week or (in aliquots) at -20°C to -80°C for longer. Avoid

repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

RefSeq: NP 001139663

 Locus ID:
 9019

 UniProt ID:
 095297

 Cytogenetics:
 1q24.2

Synonyms: MPZL1b; PZR; PZR1b; PZRa; PZRb





Summary: Cell surface receptor, which is involved in signal transduction processes. Recruits

PTPN11/SHP-2 to the cell membrane and is a putative substrate of PTPN11/SHP-2. Is a major receptor for concanavalin-A (ConA) and is involved in cellular signaling induced by ConA, which probably includes Src family tyrosine-protein kinases. Isoform 3 seems to have a dominant negative role; it blocks tyrosine phosphorylation of MPZL1 induced by ConA. Isoform 1, but not isoform 2 and isoform 3, may be involved in regulation of integrin-

mediated cell motility.[UniProtKB/Swiss-Prot Function]

Protein Families: Druggable Genome, Transmembrane

Protein Pathways: Cell adhesion molecules (CAMs)

Product images:

