

## **Product datasheet for AR51254PU-S**

## Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

EU: info-de@origene.com CN: techsupport@origene.cn

## SIRT5 (34-310, His-tag) Human Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** SIRT5 (34-310, His-tag) human recombinant protein, 0.1 mg

Species: Human
Expression Host: E. coli

**Expression cDNA Clone** MGSSHHHHHH SSGLVPRGSH MGSARPSSSM ADFRKFFAKA KHIVIISGAG VSAESGVPTF

or AA Sequence: RGAGGYWRKW QAQDLATPLA FAHNPSRVWE FYHYRREVMG SKEPNAGHRA IAECETRLGK

QGRRVVVITQ NIDELHRKAG TKNLLEIHGS LFKTRCTSCG VVAENYKSPI CPALSGKGAP EPGTQDASIP

VEKLPRCEEA GCGGLLRPHV VWFGENLDPA ILEEVDRELA HCDLCLVVGT SSVVYPAAMF

APQVAARGVP VAEFNTETTP ATNRFRFHFQ GPCGTTLPEA LACHENETVS

Tag: His-tag
Predicted MW: 32.5 kDa
Concentration: lot specific

Purity: >90% by SDS - PAGE

**Buffer:** Presentation State: Purified

State: Liquid purified protein

Buffer System: 20 mM Tris-HCl buffer (pH 8.0) containing 0.15M NaCl, 30% glycerol, 1 mM

DTT

**Preparation:** Liquid purified protein

**Protein Description:** Recombinant human SIRT5 protein, fused to His-tag at N-terminus, was expressed in E.coli

and purified by using conventional chromatography techniques.

Storage: Store undiluted at 2-8°C for one week or (in aliquots) at -20°C to -80°C for longer. Avoid

repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

**RefSeq:** NP 001180196

Locus ID:23408UniProt ID:Q9NXA8Cytogenetics:6p23Synonyms:SIR2L5





**Summary:** 

This gene encodes a member of the sirtuin family of proteins, homologs to the yeast Sir2 protein. Members of the sirtuin family are characterized by a sirtuin core domain and grouped into four classes. The functions of human sirtuins have not yet been determined; however, yeast sirtuin proteins are known to regulate epigenetic gene silencing and suppress recombination of rDNA. Studies suggest that the human sirtuins may function as intracellular regulatory proteins with mono-ADP-ribosyltransferase activity. The protein encoded by this gene is included in class III of the sirtuin family. Alternative splicing of this gene results in multiple transcript variants. [provided by RefSeq, Jul 2010]

**Protein Families:** 

Druggable Genome, Transcription Factors

## **Product images:**

