

## Product datasheet for **AR51479PU-N**

### CPEB1 (1-561, His-tag) Human Protein

#### Product data:

<b>Product Type:</b>	Recombinant Proteins
<b>Description:</b>	CPEB1 (1-561, His-tag) human recombinant protein, 0.5 mg
<b>Species:</b>	Human
<b>Expression Host:</b>	E. coli
<b>Expression cDNA Clone or AA Sequence:</b>	MGSSHHHHHH SSGLVPRGSH MGSMAFPLEE EAGRIKDCWD NQEAPALSTC SNANIFRRIN AILDNSLDFS RVCTTPINRG IHDHLPDFQD SEETVTSRML FPTSAQESSR GLPDANDLCL GLQSLSLTGW DRPWSTQDSD SSAQSSTHSV LSMLHNPLGN VLGKPPSLFL PLDPLGSDLV DKFPAPSVRG SRLDTRPILD SRSSSPSDSD TSGFSSGSDH LSDLISSLRI SPPLPFLSLS GGGPRDPLKM GVGSRMDQEQ AALAAVTPSP TSASKRWPGA SVWPSWDLLE APKDPFSIER EARLHRQAAA VNEATCTWSG QLPPRNYKNP IYSCKVFLGG VPWDITEAGL VNTFRVFGSL SVEWPGKDGK HPRCPPKGYV YLVFELEKSV RSLAQCSHD PLSPDGLSEY YFKMSSRRMR CKEVQVIPWV LADSNFVRSP SQRLDPSRTV FVGALHGMLN AEALAAAILND LFGGVVYAGI DTDKHKYPIG SGRVTFNNQR SYLKAVSAAF VEIKTTKFTK KVQIDPYLED SLCHICSSQP GPFPCRQDQC FKYFCRSCWH WRHSMEGLRH HSPLMRNQKN RDSS
<b>Tag:</b>	His-tag
<b>Predicted MW:</b>	64.5 kDa
<b>Concentration:</b>	lot specific
<b>Purity:</b>	>85% by SDS - PAGE
<b>Buffer:</b>	Presentation State: Purified State: Liquid purified protein Buffer System: 20 mM Tris-HCl buffer (pH 8.0) containing 0.4M Urea, 10% glycerol
<b>Preparation:</b>	Liquid purified protein
<b>Protein Description:</b>	Recombinant human CPEB1 protein, fused to His-tag at N-terminus, was expressed in E.coli.
<b>Storage:</b>	Store undiluted at 2-8°C for one week or (in aliquots) at -20°C to -80°C for longer. Avoid repeated freezing and thawing.
<b>Stability:</b>	Shelf life: one year from despatch.
<b>RefSeq:</b>	<u><a href="#">NP_001073001</a></u>
<b>Locus ID:</b>	64506



[View online »](#)

UniProt ID:	<a href="#">Q9BZB8</a>
Cytogenetics:	15q25.2
Synonyms:	CPE-BP1; CPEB; CPEB-1; h-CPEB; hCPEB-1
Summary:	This gene encodes a member of the cytoplasmic polyadenylation element binding protein family. This highly conserved protein binds to a specific RNA sequence, called the cytoplasmic polyadenylation element, found in the 3' untranslated region of some mRNAs. The encoded protein functions in both the cytoplasm and the nucleus. It is involved in the regulation of mRNA translation, as well as processing of the 3' untranslated region, and may play a role in cell proliferation and tumorigenesis. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2014]
Protein Pathways:	Dorso-ventral axis formation, Oocyte meiosis, Progesterone-mediated oocyte maturation

### Product images:

