

Product datasheet for **AR51531PU-N**

IDO1 / INDO (1-403, His-tag) Human Protein

Product data:

Product Type:	Recombinant Proteins
Description:	IDO1 / INDO (1-403, His-tag) human recombinant protein, 0.5 mg
Species:	Human
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	MGSSHHHHHH SSGLVPRGSH MGSMAMAHAMEN SWTISKEYHI DEEVGFALPN PQENLPDFYN DWMFIAKHLP DLIESGQLRE RVEKLNMLSI DHLTDHKSQR LARLVGCIT MAYVWGKGHG DVRKVLPRNI AVPYCQLSKK LELPPILVYA DCVLANWKKK DPNKPLTYEN MDVLFSDRDG DCSKGFFLVS LLVEIAAASA IKVIPTVFKA MQMQRDTLL KALLEIASCL EKALQVFHQI HDHVNPKAFF SVLRIYLSGW KGNPQLSDGL VYEGFWEDPK EFAGGSAGQS SVFQCFDVLG GIQQTAGGGH AAQFLQDMRR YMPPAHRNFL CSLESNPSVR EFVLSKGDAG LREAYDACVK ALVSLRSYHL QIVTKYILIP ASQQPKENKT SEDPSKLEAK GTGGTDL MNF LKTVRSTTEK SLLKEG
Tag:	His-tag
Predicted MW:	47.7 kDa
Concentration:	lot specific
Purity:	>90% by SDS - PAGE
Buffer:	Presentation State: Purified State: Liquid purified protein Buffer System: 20 mM Tris-HCl buffer (pH 8.0) containing 0.15M NaCl, 20% glycerol, 1 mM DTT
Preparation:	Liquid purified protein
Protein Description:	Recombinant human IDO1 protein, fused to His-tag at N-terminus, was expressed in E.coli and purified by using conventional chromatography techniques.
Storage:	Store undiluted at 2-8°C for one week or (in aliquots) at -20°C to -80°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
RefSeq:	NP_002155
Locus ID:	3620
UniProt ID:	P14902 , A0A348GSI3



[View online »](#)

Cytogenetics: 8p11.21

Synonyms: IDO; IDO-1; INDO

Summary: This gene encodes indoleamine 2,3-dioxygenase (IDO) - a heme enzyme that catalyzes the first and rate-limiting step in tryptophan catabolism to N-formyl-kynurenine. This enzyme acts on multiple tryptophan substrates including D-tryptophan, L-tryptophan, 5-hydroxy-tryptophan, tryptamine, and serotonin. This enzyme is thought to play a role in a variety of pathophysiological processes such as antimicrobial and antitumor defense, neuropathology, immunoregulation, and antioxidant activity. Through its expression in dendritic cells, monocytes, and macrophages this enzyme modulates T-cell behavior by its peri-cellular catabolization of the essential amino acid tryptophan.[provided by RefSeq, Feb 2011]

Protein Families: Druggable Genome

Protein Pathways: Metabolic pathways, Tryptophan metabolism

Product images:

