

Product datasheet for AR51697PU-S

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CDK1 (1-297, His-tag) Human Protein

Product data:

Product Type: Recombinant Proteins

Description: CDK1 (1-297, His-tag) human recombinant protein, 50 µg

Species: Human

Expression cDNA Clone

MEDYTKIEKI GEGTYGVVYK GRHKTTGQVV AMKKIRLESE EEGVPSTAIR EISLLKELRH PNIVSLQDVL or AA Sequence:

MQDSRLYLIF EFLSMDLKKY LDSIPPGQYM DSSLVKSYLY QILQGIVFCH SRRVLHRDLK

PQNLLIDDKG TIKLADFGLA RAFGIPIRVY THEVVTLWYR SPEVLLGSAR YSTPVDIWSI GTIFAELATK

KPLFHGDSEI DQLFRIFRAL GTPNNEVWPE VESLQDYKNT FPKWKPGSLA SHVKNLDENG

LDLLSKMLIY DPAKRISGKM ALNHPYFNDL DNQIKKMLEH HHHHH

Tag: His-tag Predicted MW: 35.1 kDa

Concentration: lot specific

Purity: >95% by SDS - PAGE

Buffer: Presentation State: Purified

State: Liquid purified protein

Buffer System: Phosphate buffered saline (pH 7.4), 30% glycerol, 2 mM DTT, 0.1 mM PMSF

Liquid purified protein Preparation:

Protein Description: Recombinant human CDK1 protein, fused to His-tag at C-terminus, was expressed in insect

cells using baculovirus expression system and purified by using conventional

chromatography techniques.

Store undiluted at 2-8°C for one week or (in aliquots) at -20°C to -80°C for longer. Avoid Storage:

repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

RefSeq: NP 001163877

983 Locus ID:

UniProt ID: B7Z3D6 Cytogenetics: 10q21.2

Synonyms: CDK1, CDC2, CDC28A, CDKN1, P34CDC2, p34 protein kinase





Summary:

The protein encoded by this gene is a member of the Ser/Thr protein kinase family. This protein is a catalytic subunit of the highly conserved protein kinase complex known as M-phase promoting factor (MPF), which is essential for G1/S and G2/M phase transitions of eukaryotic cell cycle. Mitotic cyclins stably associate with this protein and function as regulatory subunits. The kinase activity of this protein is controlled by cyclin accumulation and destruction through the cell cycle. The phosphorylation and dephosphorylation of this protein also play important regulatory roles in cell cycle control. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Mar 2009]


Protein Families:

Druggable Genome, Protein Kinase, Stem cell - Pluripotency

Protein Pathways:

Cell cycle, Gap junction, Oocyte meiosis, p53 signaling pathway, Progesterone-mediated oocyte maturation

Product images:



15% SDS-PAGE (3ug)