

Product datasheet for **AR51720PU-S**

VEGF-A (206-325, His-tag) Rat Protein

Product data:

Product Type:	Recombinant Proteins
Description:	VEGF-A (206-325, His-tag) rat recombinant protein, 20 µg
Species:	Rat
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	MGSSHHHHHH SSGLVPRGSH MGSHPAPTTE GEQKAHEVVK FMDVYQRSYC RPIETLVDIF QEYPDEIEYI FKPSCVPLMR CAGCCNDEAL ECVPTSESNV TMQIMRIKPH QSQHIGEMSF LQHSRCECRP KKDRTKPEKC DKPRR
Tag:	His-tag
Predicted MW:	16.7 kDa
Concentration:	lot specific
Purity:	>90% by SDS - PAGE
Buffer:	Presentation State: Purified State: Liquid purified protein Buffer System: Phosphate buffer saline (pH 7.4) containing 50% glycerol.
Preparation:	Liquid purified protein
Protein Description:	Recombinant Rat VEGFA protein, fused to His-tag at N-terminus, was expressed in E.coli and purified by using conventional chromatography techniques.
Storage:	Store undiluted at 2-8°C for one week or (in aliquots) at -20°C to -80°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
RefSeq:	NP_001103803
Locus ID:	83785
UniProt ID:	B5DEK7
Cytogenetics:	9q12
Synonyms:	Vegf; VEGF-A; VEGF111; VEGF164; VPF



[View online »](#)

Summary:

This gene is a member of the PDGF/VEGF growth factor family. It encodes a heparin-binding protein, which exists as a disulfide-linked homodimer. This growth factor induces proliferation and migration of vascular endothelial cells, and is essential for both physiological and pathological angiogenesis. Disruption of this gene in mice resulted in abnormal embryonic blood vessel formation. This gene is upregulated in many known tumors and its expression is correlated with tumor stage and progression. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. There is also evidence for alternative translation initiation from upstream non-AUG (CUG) codons resulting in additional isoforms. A recent study showed that a C-terminally extended isoform is produced by use of an alternative in-frame translation termination codon via a stop codon readthrough mechanism, and that this isoform is antiangiogenic. Expression of some isoforms derived from the AUG start codon is regulated by a small upstream open reading frame, which is located within an internal ribosome entry site. [provided by RefSeq, Nov 2015]

Product images: