

## Product datasheet for **AR51767PU-S**

### ATP6V1F (1-119, His-tag) Human Protein

#### Product data:

|                                       |   |
|---------------------------------------|---|
| Product Type:                         | Recombinant Proteins  |
| Description:                          | ATP6V1F (1-119, His-tag) human recombinant protein, 0.1 mg  |
| Species:                              | Human   |
| Expression Host:                      | E. coli   |
| Expression cDNA Clone or AA Sequence: | MGSSHHHHHH SSGLVPRGSH MGSMAGRGKL IAVIGDEDTV TGFLGGIGE LNKNRHPNFL VVEKDTTINE IEDTFRQFLN RDDIGIILIN QYIAEMVRHA LDAHQQSIPA VLEIPSKEHP YDAAKDSILR RARGMFTAED LR |
| Tag:                                  | His-tag   |
| Predicted MW:                         | 15.8 kDa  |
| Concentration:                        | lot specific  |
| Purity:                               | >90% by SDS - PAGE  |
| Buffer:                               | Presentation State: Purified<br>State: Liquid purified protein<br>Buffer System: Phosphate buffered saline (pH 7.4) containing, 50% glycerol, 1 mM DTT      |
| Preparation:                          | Liquid purified protein   |
| Protein Description:                  | Recombinant human ATP6V1F, fused to His-tag at N-terminus, was expressed in E.coli and purified by using conventional chromatography techniques.            |
| Storage:                              | Store undiluted at 2-8°C for one week or (in aliquots) at -20°C to -80°C for longer.<br>Avoid repeated freezing and thawing.                                |
| Stability:                            | Shelf life: one year from despatch.   |
| RefSeq:                               | <a href="#">NP_001185838</a>  |
| Locus ID:                             | 9296  |
| UniProt ID:                           | <a href="#">Q16864</a>  |
| Cytogenetics:                         | 7q32.1  |
| Synonyms:                             | ATP6S14; VATF; Vma7   |




[View online »](#)

**Summary:**

This gene encodes a component of vacuolar ATPase (V-ATPase), a multisubunit enzyme that mediates acidification of eukaryotic intracellular organelles. V-ATPase dependent organelle acidification is necessary for such intracellular processes as protein sorting, zymogen activation, receptor-mediated endocytosis, and synaptic vesicle proton gradient generation. V-ATPase is composed of a cytosolic V1 domain and a transmembrane V0 domain. The V1 domain consists of three A and three B subunits, two G subunits plus the C, D, E, F, and H subunits. The V1 domain contains the ATP catalytic site. The V0 domain consists of five different subunits: a, c, c', c'', and d. Additional isoforms of many of the V1 and V0 subunit proteins are encoded by multiple genes or alternatively spliced transcript variants. This encoded protein is the V1 domain F subunit protein. [provided by RefSeq, Jul 2008]

**Protein Pathways:**

Epithelial cell signaling in Helicobacter pylori infection, Metabolic pathways, Oxidative phosphorylation, Vibrio cholerae infection

**Product images:**

15% SDS-PAGE (3ug)