

# **Product datasheet for AR51817PU-S**

#### **OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

## **TECK / CCL25 (His-tag) Human Protein**

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** TECK / CCL25 (His-tag) human recombinant protein, 0.1 mg

Species: Human
Expression Host: E. coli

**Expression cDNA Clone** 

or AA Sequence:

MGSSHHHHHH SSGLVPRGSH MGSHMQGVFE DCCLAYHYPI GWAVLRRAWT YRIQEVSGSC NLPAAIFYLP KRHRKVCGNP KSREVQRAMK LLDARNKVFA KLHHNTQTFQ AGPHAVKKLS

SGNSKLSSSK FSNPISSSKR NVSLLISANS GL

Tag: His-tag
Predicted MW: 16.8 kDa
Concentration: lot specific

Purity: >90% by SDS - PAGE

**Buffer:** Presentation State: Purified

State: Liquid purified protein

Buffer System: 20 mM Tris-HCl buffer (pH 8.0) containing 10% glycerol.

**Preparation:** Liquid purified protein

**Protein Description:** Recombinant protein shows chemotactic activity on thymocytes, macrophages, THP-1 cells,

and dendritics cells but is inactive on peripheral blood lymphocytes and neutrophils. It binds to CCR9. Isoform 2 is an antagonist of isoform1 and binds to atypical chemokine receptor ACKR4 and mediates the recruitment of beta-arrestin (ARRB1/2) to ACKR4. Recombinant

human CCL25-iso1, fused to His-tag at N-terminus, was expressed in E.coli.

Storage: Store undiluted at 2-8°C for one week or (in aliquots) at -20°C to -80°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

**RefSeq:** NP 001188288

 Locus ID:
 6370

 UniProt ID:
 015444

 Cytogenetics:
 19p13.2

Synonyms: Ckb15; Ck beta-15; SCYA25; TECK



#### TECK / CCL25 (His-tag) Human Protein - AR51817PU-S

**Summary:** This antimicrobial gene belongs to the subfamily of small cytokine CC genes. Cytokines are a

family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by

this gene displays chemotactic activity for dendritic cells, thymocytes, and activated


macrophages but is inactive on peripheral blood lymphocytes and neutrophils. The product of this gene binds to chemokine receptor CCR9. Alternative splicing results in multiple

transcript variants. [provided by RefSeq, Sep 2014]

**Protein Families:** Druggable Genome, Secreted Protein

**Protein Pathways:** Chemokine signaling pathway, Cytokine-cytokine receptor interaction

### **Product images:**



15% SDS-PAGE (3ug)