

## Product datasheet for AR51857PU-S

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Sonic hedgehog (SHH) (24-197, ) Human Protein

**Product data:** 

**Product Type: Recombinant Proteins** 

**Description:** Sonic hedgehog (SHH) (24-197, ) human recombinant protein, 50 μg

Species: Human E. coli **Expression Host:** 

**Expression cDNA Clone** 

MIIGPGRGFG KRRHPKKLTP LAYKQFIPNV AEKTLGASGR YEGKISRNSE RFKELTPNYN PDIIFKDEEN or AA Sequence:

TGADRLMTQR CKDKLNALAI SVMNQWPGVK LRVTEGWDED GHHSEESLHY EGRAVDITTS

DRDRSKYGML ARLAVEAGFD WVYYESKAHI HCSVKAENSV AAKSGG

Predicted MW: 19.8 kDa Concentration: lot specific

**Purity:** >95% by SDS - PAGE

**Buffer:** Presentation State: Purified

State: Liquid purified protein

Buffer System: Phosphate buffered saline (pH 7.4)

**Endotoxin:** < 1.0 Eu per 1 microgram of protein (determined by LAL method)

**Preparation:** Liquid purified protein

**Protein Description:** Recombinant human SHH was expressed in E.coli and purified by using conventional

chromatography techniques.

Store undiluted at 2-8°C for one week or (in aliquots) at -20°C to -80°C for longer. Storage:

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

RefSeq: NP 000184

Locus ID: 6469 **UniProt ID:** Q15465 Cytogenetics: 7q36.3

Synonyms: Sonic hedgehog protein, HHG-1





Summary:

This gene encodes a protein that is instrumental in patterning the early embryo. It has been implicated as the key inductive signal in patterning of the ventral neural tube, the anteriorposterior limb axis, and the ventral somites. Of three human proteins showing sequence and functional similarity to the sonic hedgehog protein of Drosophila, this protein is the most similar. The protein is made as a precursor that is autocatalytically cleaved; the N-terminal portion is soluble and contains the signalling activity while the C-terminal portion is involved in precursor processing. More importantly, the C-terminal product covalently attaches a cholesterol moiety to the N-terminal product, restricting the N-terminal product to the cell surface and preventing it from freely diffusing throughout the developing embryo. Defects in this protein or in its signalling pathway are a cause of holoprosencephaly (HPE), a disorder in which the developing forebrain fails to correctly separate into right and left hemispheres. HPE is manifested by facial deformities. It is also thought that mutations in this gene or in its signalling pathway may be responsible for VACTERL syndrome, which is characterized by vertebral defects, anal atresia, tracheoesophageal fistula with esophageal atresia, radial and renal dysplasia, cardiac anomalies, and limb abnormalities. Additionally, mutations in a long range enhancer located approximately 1 megabase upstream of this gene disrupt limb patterning and can result in preaxial polydactyly. [provided by RefSeq, Jul 2008]

Protein Families: Protein Pathways: Druggable Genome, ES Cell Differentiation/IPS, Secreted Protein, Transmembrane Basal cell carcinoma, Hedgehog signaling pathway, Pathways in cancer

## **Product images:**

