

Product datasheet for AR51863PU-S

PAFAH2 (1-392, His-tag) Human Protein

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

| Product Type: | Recombinant Proteins | |
|--|--|--|
| Description: | PAFAH2 (1-392, His-tag) human protein, 0.1 mg | |
| Species: | Human | |
| Expression Host: | E. coli | |
| Expression cDNA Clone or AA Sequence: | MGSSHHHHHH SSGLVPRGSH MGSMGVNQSV GFPPVTGPHL VGCGDVMEGQ NLQGSFFRLF YPCQKAEETM EQPLWIPRYE YCTGLAEYLQ FNKRCGGLLF NLAVGSCRLP VSWNGPFKTK DSGYPLIIFS HGLGAFRTLY SAFCMELASR GFVVAVPEHR DRSAATTYFC KQAPEENQPT NESLQEEWIP FRRVEEGEKE FHVRNPQVHQ RVSECLRVLK ILQEVTAGQT VFNILPGGLD LMTLKGNIDM SRVAVMGHSF GGATAILALA KETQFRCAVA LDAWMFPLER DFYPKARGPV FFINTEKFQT MESVNLMKKI CAQHEQSRII TVLGSVHRSQ TDFAFVTGNL IGKFFSTETR GSLDPYEGQE VMVRAMLAFL QKHLDLKEDY NQWNNLIEGI GPSLTPGAPH HLSSL | |
| Tag: | His-tag | |
| Predicted MW: | 46.4 kDa | |
| Concentration: | lot specific | |
| Purity: | >90% by SDS - PAGE | |
| Buffer: | Presentation State: Purified State: Liquid purified protein Buffer System: Phosphate buffered saline (pH 7.4) containing 20% glycerol, 1 mM DTT | |
| Preparation: | Liquid purified protein | |
| Storage: | Store undiluted at 2-8°C for one week or (in aliquots) at -20°C to -80°C for longer. Avoid repeated freezing and thawing. | |
| Stability: | Shelf life: one year from despatch. | |
| RefSeq: | <u>NP 000428</u> | |
| Locus ID: | 5051 | |
| UniProt ID: | <u>Q99487</u> | |
| Cytogenetics: | 1p36.11 | |
| Synonyms: | HSD-PLA2 | |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

| | PAFAH2 (1-392, His-tag) Human Protein – AR51863PU-S | |
|-------------------|--|--|
| Summary: | This gene encodes platelet-activating factor acetylhydrolase isoform 2, a single-subunit intracellular enzyme that catalyzes the removal of the acetyl group at the SN-2 position of platelet-activating factor (identified as 1-O-alkyl-2-acetyl-sn-glyceryl-3-phosphorylcholine). However, this lipase exhibits a broader substrate specificity than simply platelet activating factor. Two other isoforms of intracellular platelet-activating factor acetylhydrolase exist, and both are multi-subunit enzymes. Additionally, there is a single-subunit serum isoform of this enzyme. [provided by RefSeq, Jul 2008] | |
| Protein Families: | Druggable Genome | |
| Protein Pathway | Ether lipid metabolism, Metabolic pathways | |

Product images:

| (kDa) 150 100 70 | = |
|---------------------------|---|
| 50 | - |
| 35 | |
| 25 | |
| 20 | - |
| 15 | - |

15% SDS-PAGE (3ug)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US