

Product datasheet for **AR51902PU-N**

CD212 / IL12RB1 (24-545, His-tag) Human Protein

Product data:

| | |
|--|---|
| Product Type: | Recombinant Proteins |
| Description: | CD212 / IL12RB1 (24-545, His-tag) human recombinant protein, 0.25 mg |
| Species: | Human |
| Expression cDNA Clone or AA Sequence: | CRTSECCFQD PPYPDADSGS ASGPRDLRCY RISSDRYECS WQYEGPTAGV SHFLRCCLSS GRCCYFAAGS ATRLQFSDQA GSVLYTVTL WVESWARNQT EKSPEVTLQL YNSVKYEPPL GDKVSKLAG QLRMEWETPD NQVGAEVQFR HRTSPSPWKL GDCGPQDDDT ESCLCPLMN VAQEFQLRRR QLGSGSSWS KWSSPVCVPP ENPPQPQVRF SVEQLGQDGR RRLTLKEQPT QLELPEGCQG LAPGTEVTYR LQLHMLSCPC KAKATRTLHL GKMPYLSGAA YNVAVISSNQ FGPGLNQTWH IPADTHTEPV ALNISVGTNG TTMYPARAQ SMTYCIWQP VGQDGGGLATC SLTAPQDPDP AGMATYSWSR ESGAMGQEK YITIFASAH PEKLTWSTV LSTYHFGGNA SAAGTPHHVS VKNHSLDSVS VDWAPSLST CPGVLKEYVW RCRDEDSKQV SEHPVQPTET QVTLSGLRAG VAYTVQVRAD TAWLRGVWSQ PQRFSIEVQV SDHHHHHH |
| Tag: | His-tag |
| Predicted MW: | 58.4 kDa |
| Concentration: | lot specific |
| Purity: | >90% by SDS - PAGE. |
| Buffer: | Presentation State: Purified State: Liquid purified protein Buffer System: Phosphate buffered saline (pH 7.4) containing 20% glycerol, 1 mM EDTA, 0.1 mM PMSF |
| Endotoxin: | < 1.0 EU per 1 microgram of protein (determined by LAL method) |
| Preparation: | Liquid purified protein |
| Protein Description: | Recombinant human IL12RB1, fused to His-tag at C-terminus, was expressed in insect cell and purified by using conventional chromatography techniques. |
| Storage: | Store undiluted at 2-8°C for one week or (in aliquots) at -20°C to -80°C for longer. Avoid repeated freezing and thawing. |
| Stability: | Shelf life: one year from despatch. |
| RefSeq: | NP_001276952 |
| Locus ID: | 3594 |



[View online »](#)

UniProt ID: [P42701](#)

Cytogenetics: 19p13.11

Synonyms: CD212; IL-12R-BETA1; IL12RB; IMD30

Summary: The protein encoded by this gene is a type I transmembrane protein that belongs to the hemopoietin receptor superfamily. This protein binds to interleukine 12 (IL12) with a low affinity, and is thought to be a part of IL12 receptor complex. This protein forms a disulfide-linked oligomer, which is required for its IL12 binding activity. The coexpression of this and IL12RB2 proteins was shown to lead to the formation of high-affinity IL12 binding sites and reconstitution of IL12 dependent signaling. Mutations in this gene impair the development of interleukin-17-producing T lymphocytes and result in increased susceptibility to mycobacterial and Salmonella infections. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Feb 2014]

Protein Families: Druggable Genome, Transmembrane

Protein Pathways: Cytokine-cytokine receptor interaction, Jak-STAT signaling pathway

Product images:

