

Product datasheet for AR51918PU-S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Complement factor D (21-253, His-tag) Human Protein

Product data:

Product Type: Recombinant Proteins

Description: Complement factor D (21-253, His-tag) human recombinant protein, 50 μg

Species: Human

Expression cDNA Clone

or AA Sequence:

PPRGRILGGR EAEAHARPYM ASVQLNGAHL CGGVLVAEQW VLSAAHCLED AADGKVQVLL GAHSLSQPEP SKRLYDVLRA VPHPDSQPDT IDHDLLLLQL SEKATLGPAV RPLPWQRVDR DVAPGTLCDV AGWGIVNHAG RRPDSLQHVL LPVLDRATCN RRTHHDGAIT ERLMCAESNR RDSCKGDSGG PLVCGGVLEG VVTSGSRVCG NRKKPGIYTR VASYAAWIDS VLAVEHHHHH H

Tag: His-tag

Predicted MW: 26.01 kDa

Concentration: lot specific

Purity: >95% by SDS - PAGE.

Buffer: Presentation State: Purified

State: Liquid purified protein

Buffer System: Phosphate buffered saline (pH 7.4)

Endotoxin: < 1.0 Eu per 1 microgram of protein (determined by LAL method)

Preparation: Liquid purified protein

Protein Description: Recombinant human CFD, fused to His-tag at C-terminus, was expressed in insect cell and

purified by using conventional chromatography techniques.

Storage: Store undiluted at 2-8°C for one week or (in aliquots) at -20°C to -80°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

RefSeg: NP 001304264

 Locus ID:
 1675

 UniProt ID:
 P00746

 Cytogenetics:
 19p13.3

Synonyms: Properdin factor D, Adipsin, CFD, DF, PFD





Summary:

This gene encodes a member of the S1, or chymotrypsin, family of serine peptidases. This protease catalyzes the cleavage of factor B, the rate-limiting step of the alternative pathway of complement activation. This protein also functions as an adipokine, a cell signaling protein secreted by adipocytes, which regulates insulin secretion in mice. Mutations in this gene underlie complement factor D deficiency, which is associated with recurrent bacterial meningitis infections in human patients. Alternative splicing of this gene results in multiple transcript variants. At least one of these variants encodes a preproprotein that is proteolytically processed to generate the mature protease. [provided by RefSeq, Nov 2015]

Protein Families: Druggable Genome, Protease, Secreted Protein

Protein Pathways: Complement and coagulation cascades

Product images:

