

Product datasheet for AR51942PU-N

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

Cystatin-D (26-142, His-tag) Human Protein

Product data:

Product Type: Recombinant Proteins

Description: Cystatin-D (26-142, His-tag) human recombinant protein, 0.25 mg

Species: Human

Expression cDNA Clone

or AA Sequence:

QSRTLAGGIH ATDLNDKSVQ RALDFAISEY NKVINKDEYY SRPLQVMAAY QQIVGGVNYY FNVKFGRTTC TKSQPNLDNC PFNDQPKLKE EEFCSFQINE VPWEDKISIL NYKCRKVHHH HHH

Tag: His-tag

Predicted MW: 14.36 kDa

Concentration: lot specific

Purity: >90% by SDS - PAGE

Buffer: Presentation State: Purified

State: Liquid purified protein

Buffer System: Phosphate buffered saline (pH 7.4) containing 10% glycerol

Endotoxin: < 1.0 EU per 1 microgram of protein (determined by LAL method)

Preparation: Liquid purified protein

Protein Description: Recombinant human CST5, fused to His-tag at C-terminus, was expressed in insect cell and

purified by using conventional chromatography techniques.

Storage: Store undiluted at 2-8°C for one week or (in aliquots) at -20°C to -80°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

RefSeq: NP 001891

Locus ID: 1473

UniProt ID: P28325

Cytogenetics: 20p11.21

Synonyms: CST5, Cystatin-5





Summary:

The cystatin superfamily encompasses proteins that contain multiple cystatin-like sequences. Some of the members are active cysteine protease inhibitors, while others have lost or perhaps never acquired this inhibitory activity. There are three inhibitory families in the superfamily, including the type 1 cystatins (stefins), type 2 cystatins and the kininogens. The type 2 cystatin proteins are a class of cysteine proteinase inhibitors found in a variety of human fluids and secretions. The cystatin locus on chromosome 20 contains the majority of the type 2 cystatin genes and pseudogenes. This gene is located in the cystatin locus and encodes a protein found in saliva and tears. The encoded protein may play a protective role against proteinases present in the oral cavity. [provided by RefSeq, Jul 2008]

Product images:

