

# **Product datasheet for AR51978PU-N**

#### 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

## Cystatin-S (21-141, His-tag) Human Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Cystatin-S (21-141, His-tag) human protein, 0.25 mg

Species: Human
Expression Host: Insect

**Expression cDNA Clone** ADPMSSSKEE NRIIPGGIYD ADLNDEWVQR ALHFAISEYN KATEDEYYRR PLQVLRAREQ

or AA Sequence: TFGGVNYFFD VEVGRTICTK SQPNLDTCAF HEQPELQKKQ LCSFEIYEVP WEDRMSLVNS

RCQEAHHHHH H

Tag: His-tag

Predicted MW: 15.4 kDa

Concentration: lot specific

Purity: >95% by SDS - PAGE

**Buffer:** Presentation State: Purified

State: Liquid purified protein

Buffer System: Phosphate Buffered Saline (pH 7.4) containing 10% glycerol.

**Endotoxin:** < 1.0 EU per 1 microgram of protein (determined by LAL method)

**Preparation:** Liquid purified protein

Storage: Store undiluted at 2-8°C for one week or (in aliquots) at -20°C to -80°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

**RefSeq:** <u>NP 001890</u>

 Locus ID:
 1472

 UniProt ID:
 P01036

 Cytogenetics:
 20p11.21






#### **Summary:**

The cystatin superfamily encompasses proteins that contain multiple cystatin-like sequences. Some of the members are active cysteine protease inhibitors, while others have lost or perhaps never acquired this inhibitory activity. There are three inhibitory families in the superfamily, including the type 1 cystatins (stefins), type 2 cystatins and the kininogens. The type 2 cystatin proteins are a class of cysteine proteinase inhibitors found in a variety of human fluids and secretions. The cystatin locus on chromosome 20 contains the majority of the type 2 cystatin genes and pseudogenes. This gene is located in the cystatin locus and encodes a type 2 salivary cysteine peptidase inhibitor. The protein is an S-type cystatin, based on its high level of expression in saliva, tears and seminal plasma. The specific role in these fluids is unclear but antibacterial and antiviral activity is present, consistent with a protective function. [provided by RefSeq, Jul 2008]

## **Product images:**



15% SDS-PAGE (3ug)