

## Product datasheet for AR51993PU-N

## Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

## PGD / PGDH (1-483, His-tag) Human Protein

**Product data:** 

**Product Type: Recombinant Proteins** 

**Description:** PGD / PGDH (1-483, His-tag) human protein, 0.1 mg

Species: Human **Expression Host:** E. coli

**Expression cDNA Clone** 

MGSSHHHHHH SSGLVPRGSH MAQADIALIG LAVMGQNLIL NMNDHGFVVC AFNRTVSKVD or AA Sequence:

DFLANEAKGT KVVGAQSLKE MVSKLKKPRR IILLVKAGQA VDDFIEKLVP LLDTGDIIID GGNSEYRDTT

RRCRDLKAKG ILFVGSGVSG GEEGARYGPS LMPGGNKEAW PHIKTIFQGI AAKVGTGEPC

CDWVGDEGAG HFVKMVHNGI EYGDMQLICE AYHLMKDVLG MAQDEMAQAF EDWNKTELDS FLIEITANIL KFQDTDGKHL LPKIRDSAGQ KGTGKWTAIS ALEYGVPVTL IGEAVFARCL SSLKDERIQA

SKKLKGPQKF QFDGDKKSFL EDIRKALYAS KIISYAQGFM LLRQAATEFG WTLNYGGIAL MWRGGCIIRS VFLGKIKDAF DRNPELQNLL LDDFFKSAVE NCQDSWRRAV STGVQAGIPM PCFTTALSFY DGYRHEMLPA SLIQAQRDYF GAHTYELLAK PGQFIHTNWT GHGGTVSSSS YNA

Tag: His-tag Predicted MW: 55.3 kDa Concentration: lot specific

**Purity:** >90% by SDS - PAGE

**Buffer:** Presentation State: Purified

State: Liquid purified protein

Buffer System: 20 mM Tris-HCl buffer (pH 8.0) containing 1 mM DTT, 0.1M Nacl, 10% glycerol

Specific: Specific activity is > 10 units/mg, in which one unit oxidize 1.0 umole of 6-phospho-D-**Bioactivity:** 

gluconate to D-ribulose 5-phosphate per minute at pH 8.0 at 25C, in the presence of ß-NADP.

**Preparation:** Liquid purified protein

Store undiluted at 2-8°C for one week or (in aliquots) at -20°C to -80°C for longer. Storage:

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

RefSeq: NP 001291380

5226 Locus ID:

UniProt ID: P52209, B4E2U0





Cytogenetics: 1p36.22

Synonyms: 6PGD

**Summary:** 6-phosphogluconate dehydrogenase is the second dehydrogenase in the pentose phosphate

shunt. Deficiency of this enzyme is generally asymptomatic, and the inheritance of this disorder is autosomal dominant. Hemolysis results from combined deficiency of 6-

phosphogluconate dehydrogenase and 6-phosphogluconolactonase suggesting a synergism of the two enzymopathies. Several transcript variants encoding different isoforms have been

found for this gene. [provided by RefSeq, Jan 2015]

**Protein Pathways:** Glutathione metabolism, Metabolic pathways, Pentose phosphate pathway

## **Product images:**

