

Product datasheet for AR52066PU-N

CD38 (43-300, His-tag) Human Protein

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Recombinant Proteins
Description:	CD38 (43-300, His-tag) human protein, 0.25 mg
Species:	Human
Expression cDNA Clone or AA Sequence:	ADPEFVPRWR QQWSGPGTTK RFPETVLARC VKYTEIHPEM RHVDCQSVWD AFKGAFISKH PCNITEEDYQ PLMKLGTQTV PCNKILLWSR IKDLAHQFTQ VQRDMFTLED TLLGYLADDL TWCGEFNTSK INYQSCPDWR KDCSNNPVSV FWKTVSRRFA EAACDVVHVM LNGSRSKIFD KNSTFGSVEV HNLQPEKVQT LEAWVIHGGR EDSRDLCQDP TIKELESIIS KRNIQFSCKN IYRPDKFLQC VKNPEDSSCT SEIHHHHHH
Tag:	His-tag
Predicted MW:	31.2 kDa
Concentration:	lot specific
Purity:	>90% by SDS - PAGE
Buffer:	Presentation State: Purified State: Liquid purified protein Buffer System: 50 mM MES (pH 5.0) containing 100 mM NaCl, 10% glycerol
Endotoxin:	< 1.0 EU per 1 microgram of protein (determined by LAL method)
Preparation:	Liquid purified protein
Storage:	Store undiluted at 2-8°C for one week or (in aliquots) at -20°C to -80°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
RefSeq:	<u>NP 001766</u>
Locus ID:	952
UniProt ID:	<u>P28907</u> , <u>B4E006</u>
Cytogenetics:	4p15.32
Synonyms:	ADPRC 1; ADPRC1




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

CD38 (43-300, His-tag) Human Protein – AR52066PU-N

Summary:	The protein encoded by this gene is a non-lineage-restricted, type II transmembrane glycoprotein that synthesizes and hydrolyzes cyclic adenosine 5'-diphosphate-ribose, an intracellular calcium ion mobilizing messenger. The release of soluble protein and the ability of membrane-bound protein to become internalized indicate both extracellular and intracellular functions for the protein. This protein has an N-terminal cytoplasmic tail, a single membrane-spanning domain, and a C-terminal extracellular region with four N-glycosylation sites. Crystal structure analysis demonstrates that the functional molecule is a dimer, with the central portion containing the catalytic site. It is used as a prognostic marker for patients with chronic lymphocytic leukemia. Alternative splicing results in multiple transcript variants. [provided by RefSeg, Sep 2015]
Protein Families:	Cancer stem cells, Druggable Genome, ES Cell Differentiation/IPS, Induced pluripotent stem cells, Transmembrane

Protein Pathways:Calcium signaling pathway, Hematopoietic cell lineage, Metabolic pathways, Nicotinate and
nicotinamide metabolism

Product images:



15% SDS-PAGE (3ug)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US