

Product datasheet for AR52070PU-N

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

CD153 / CD30L (63-234, His-tag) Human Protein

Product data:

Product Type: Recombinant Proteins

Description: CD153 / CD30L (63-234, His-tag) human protein, 0.1 mg

Species: Human

Expression cDNA Clone

or AA Sequence:

ADPQRTDSIP NSPDNVPLKG GNCSEDLLCI LKRAPFKKSW AYLQVAKHLN KTKLSWNKDG ILHGVRYQDG NLVIQFPGLY FIICQLQFLV QCPNNSVDLK LELLINKHIK KQALVTVCES GMQTKHVYQN LSQFLLDYLQ VNTTISVNVD TFQYIDTSTF PLENVLSIFL YSNSDHHHHH H

Tag: His-tag

Predicted MW: 20.7 kDa

Concentration: lot specific

Purity: >85% by SDS - PAGE

Buffer: Presentation State: Purified

State: Liquid purified protein

Buffer System: Phosphate Buffered Saline (pH 7.4) containing 10% glycerol.

Endotoxin: < 1.0 EU per 1 microgram of protein (determined by LAL method)

Preparation: Liquid purified protein

Storage: Store undiluted at 2-8°C for one week or (in aliquots) at -20°C to -80°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

RefSeq: NP 001235

Locus ID: 944

 UniProt ID:
 P32971

 Cytogenetics:
 9q32-q33.1

Synonyms: TNFSF8, CD30 Ligand, CD30LG, CD30-L






Summary:

The protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This cytokine is a ligand for TNFRSF8/CD30, which is a cell surface antigen and a marker for Hodgkin lymphoma and related hematologic malignancies. The engagement of this cytokine expressed on B cell surface plays an inhibitory role in modulating Ig class switch. This cytokine was shown to enhance cell proliferation of some lymphoma cell lines, while to induce cell death and reduce cell proliferation of other lymphoma cell lines. The pleiotropic biologic activities of this cytokine on different CD30+ lymphoma cell lines may play a pathophysiologic role in Hodgkin's and some non-Hodgkin's lymphomas. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Nov 2011]

Protein Families: Druggable Genome, Transmembrane
Protein Pathways: Cytokine-cytokine receptor interaction

Product images:



15% SDS-PAGE (3ug)