

Product datasheet for BA1008S

Cytokeratin 18 Bovine Protein

Product data:

Product Type: Native Proteins

Description: Cytokeratin 18 bovine protein, 0.1 mg

Species: Bovine **Protein Source:** Liver Predicted MW: 45 kDa

Concentration: lot specific

Purity: >95% (determined by SDS gelelectrophoresis)

Buffer: Presentation State: Purified

State: Lyophilized

Buffer System: 30 mM Tris/HCl pH 8, 9M urea, 2 mM DTT, 2 mM EDTA, 10 mM

Methylammonium Chloride

Reconstitution Method: Restore with distilled water.

> BA1008S: 70 μl (final volume 100 μl). BA1008 : 175 μl (final volume 250 μl).

Preparation: Lyophilized

Applications: Protein standard in 1D and 2D SDS gelelectrophoresis.

> Immunoassays. Immunization.

Protocol: Reconstitution to filaments is performed by mixing equimolar amounts of keratins of type I and type II at concentrations of approx. 0.5 mg/ml, both dissolved in 9.5 M urea buffer (see above). Protofilaments and filament complexes are obtained by dialyzing the resulting polypeptide solution stepwise to a concentration of 4 M urea and then to low salt

condition (50 mM NaCl, 2 mM dithiothreitol, 10 mM Tris-HCl, pH 7.4).

For Immunization purposes, the solution can be further dialyzed against PBS (phosphate

buffered saline, e.g. Dulbeccos PBS). See References 2 and 3 for more details.

Protein Description: Bovine keratin K18 (formerly also designated Cytokeratin 18).

Note: Isoelectric Point: pl 5.4



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

ORÏGENE

Cytokeratin 18 Bovine Protein - BA1008S

Storage: Store at 2-8°C (lyophilized) and at -20°C (reconstituted).

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

RefSeq: NP 000215

Locus ID: 3875

Cytogenetics: 12q13.13

Synonyms: CK-18; CYK18; K18

Summary: KRT18 encodes the type I intermediate filament chain keratin 18. Keratin 18, together with its

filament partner keratin 8, are perhaps the most commonly found members of the

intermediate filament gene family. They are expressed in single layer epithelial tissues of the body. Mutations in this gene have been linked to cryptogenic cirrhosis. Two transcript variants encoding the same protein have been found for this gene. [provided by RefSeq, Jul

2008]

Protein Pathways: Pathogenic Escherichia coli infection

Product images:

a --

b —

c —

d —

1 2