

Product datasheet for BA1011S

GFAP Bovine Protein

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Native Proteins
21	
Description:	GFAP bovine protein, 0.1 mg
Species:	Bovine
Protein Source:	Spinal Cord
Predicted MW:	52 kDa
Concentration:	lot specific
Purity:	>98% (determined by SDS gelelectrophoresis).
Buffer:	Presentation State: Purified State: Lyophilized Buffer System: Final Solution contains 10 mM Sodium Phosphate buffer pH 7.5, 6M Urea, 2 mM DTT, 1 mM EDTA, 10 mM Methylammonium Chloride
Reconstitution Method:	Restore with distilled water. BA1011S: 80 μl (final volume 100 μl). BA1011 : 200 μl (final volume 250 μl).
Preparation:	Lyophilized
Applications:	 Protein standard in 1D and 2D SDS gelelectrophoresis. Immunoassays. Immunization. Protocol: Reconstitution to Filaments is performed by dissolving in 6 M urea buffer (see above) at concentrations of approx. 0.5 mg/ml. Protofilaments and filament complexes are obtained by dialyzing the resulting polypeptide solution stepwise to a concentration of 4 M urea and then to low salt condition (50 mM NaCl, 2 mM dithiothreitol, 10 mM Tris-HCl, pH 7.4). For immunization purposes, the solution can be further dialyzed against PBS (phosphate buffered saline, e.g. Dulbeccos PBS).
Protein Description:	Bovine Glial Filament Protein (GFP)
Note:	Isoelectric Point: pl 5.4




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	GFAP Bovine Protein – BA1011S
Storage:	Prior to reconstitution store at 2-8°C. Following reconstitution store the protein undiluted at -20°C. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
RefSeq:	<u>NP 776490</u>
Locus ID:	281189
Synonyms:	Glial Fibrillary Acidic Protein
Summary:	glial fibrillary acidic protein
Protein Families	 Protein standard in 1D and 2D SDS gelelectrophoresis. Immunoassays. Immunization. Protocol: Reconstitution to Filaments is performed by dissolving in 6 M urea buffer (see above) at concentrations of approx. 0.5 mg/ml. Protofilaments and filament complexes are obtained by dialyzing the resulting polypeptide solution stepwise to a concentration of 4 M urea and then to low salt condition (50 mM NaCl, 2 mM dithiothreitol, 10 mM Tris-HCl, pH 7.4). For immunization purposes, the solution can be further dialyzed against PBS (phosphate

For immunization purposes, the solution can be further dialyzed against PBS (phosphate buffered saline, e.g. Dulbeccos PBS).

Product images:



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US