

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **Product datasheet for BIN093**

## Hepatitis B Surface Antigen / HBsAg (adr) Protein

## **Product data:**

Product Type:	Recombinant Proteins
Description:	Hepatitis B Surface Antigen / HBsAg (adr) recombinant protein, 0.5 mg
Expression Host:	S. cerevisiae
Concentration:	lot specific
Purity:	>98% lon exchange chromatography. Product is sterile filtered.
Buffer:	Presentation State: Purified State: Liquid purified lg fraction (>95% pure by HPLC). Buffer System: 10 mM Phosphate and 145 mM Sodium Chloride as buffer, containing 0.09% Sodium Azide as preservative.
Preparation:	Liquid purified lg fraction (>95% pure by HPLC).
Applications:	Suitable for use in ELISA (capture and conjugate), Immunochromatography (capture and conjugate) and Immunogen for monoclonal antibody production. Tested in ELISA with anti HBsAg antibodies.
Protein Description:	Recombinant Hepatitis B Surface Antigen (HBsAg),(adr)(S Region). Recombinant HBsAg is a 24 kDa protein. Does not contain a fusion partner.
Storage:	Store the antigen at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid multiple freeze/thaw cycles.
Stability:	Shelf life: six months from despatch.
Synonyms:	HBV surface antigen, Hepatitis B Virus
Summary:	Hepatitis B Virus (HBV) infection induces a disease state which manifests itself in a variety of ways, characterized by the extent of liver damage, inflammation and viral persistence. HBV infection is also associated with a 100 fold increased risk of hepatocellular carcinoma and currently infects over 250 million people worldwide. HBV has a partially double stranded 3.2 kilobase DNA genome which contains four open reading frames. One of these encodes a 154 amino acid protein called the HBx protein. HBx has been shown to be a transcriptional transactivator of both viral and cellular promoters. Lacking a DNA binding domain and nuclear localization signal, HBx is believed to exert transcriptional activity through protein protein interaction.



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Protein Families:Suitable for use in ELISA (capture and conjugate), Immunochromatography (capture and<br/>conjugate) and Immunogen for monoclonal antibody production. Tested in ELISA with anti<br/>HBsAg antibodies.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US