

## Product datasheet for **BM2274**

### EPCAM Mouse Monoclonal Antibody [Clone ID: HEA125]

#### Product data:

Product Type:	Primary Antibodies
Clone Name:	HEA125
Applications:	IF, IHC, WB
Recommended Dilution:	<b>Immunofluorescence Microscopy.</b> <b>Cell Sorting.</b> <b>Cytological material.</b> <b>Immunoblotting.</b> <b>Immunohistochemistry on Frozen Sections and Paraffin-Embedded Sections.</b> With paraffin embedded sections, protease pre-treatment is required prior to antibody application. <b>Recommended Dilutions:</b> Dilute at least 1/10 with PBS, pH 7.4, for Immunohistochemical application. <b>Incubation time:</b> 1 h at RT; extended with paraffin.
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	HT-29 colon carcinoma cell line
Specificity:	The Ep-CAM (HEA125) antibody recognizes the epithelial cell adhesion molecule Ep-CAM (also described as 17-1A antigen or EPG34). This antigen is widely expressed on cells of epithelial origin and tumors derived therefrom. HEA125 represents an excellent marker to discriminate epithelial from mesothelial structures. The antigen has been detected in all carcinoma types tested (18 different origins). A subset of squamous cell carcinoma is negative. <b>Tested Reactivities on Cultured Cell Lines:</b> All carcinoma cell lines tested so far; particularly strong reaction with colon carcinoma cell lines (HT-29, WiDr, SW1116). <b>Polypeptide Reacting:</b> Mr 40,000 human epithelium-specific cell surface glycoprotein (Ep-CAM).



[View online »](#)

<b>Formulation:</b>	PBS, pH 7.4 State: Purified State: Lyophilized purified Ig fraction Stabilizer: 0.5% BSA Preservative: 0.09% Sodium Azide Label: Restore in 1.0 ml distilled water
<b>Purification:</b>	Affinity Chromatography on Protein A
<b>Conjugation:</b>	Unconjugated
<b>Storage:</b>	Prior to reconstitution store at 2-8°C. Following reconstitution store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
<b>Stability:</b>	Shelf life: one year from despatch.
<b>Gene Name:</b>	epithelial cell adhesion molecule
<b>Database Link:</b>	<a href="#">Entrez Gene 4072 Human P16422</a>
<b>Background:</b>	Epithelial Cell Adhesion Molecule (EpCAM) is a 40 kDa cell surface antigen. This antigen has been identified independently by a number of groups, and has been known by a variety of names. Several monoclonal antibodies have been raised against EpCAM, many of which have been described as tumour specific molecules on carcinomas. EpCAM is a Type 1 transmembrane glycoprotein. It is expressed on the basolateral membrane of cells by the majority of epithelial tissues, with the exception of adult squamous epithelium and some specific epithelial cell types including hepatocytes and gastric epithelial cells. EpCAM expression has been reported to be a possible marker of early malignancy, with expression being increased in tumour cells, and de novo expression being seen in dysplastic squamous epithelium.
<b>Synonyms:</b>	Ep-CAM, Epithelial cell adhesion molecule, GA733-2, EGP314, KSA, TROP1, Trop-1, M1S2, M4S1, MIC18