

Product datasheet for **BM4018S**

Fibroblasts (Pan Reticular) Rat Monoclonal Antibody [Clone ID: ER-TR7]

Product data:

Product Type: Primary Antibodies

Clone Name: ER-TR7

Applications: FC, IF, IHC

Recommended Dilution: **Immunohistochemistry on Frozen Sections** (Ref.1,2): Sections were stained using an indirect immunoperoxidase method (Ref.1).

Immunohistochemistry on Paraffin Sections: Formalin fixed paraffin sections were deparafined, hydrated in ethanol and stained with *ER-TR7* for 30' at RT (Ref.3).

Flow Cytometry: Splenocytes were incubated with *ER-TR7* for 30' (Ref.4).

Immunofluorescence (Ref.4-6): Acetone or PFA fixed cells were quenched with 50mM NH_4Cl for 30', blocked and permeabilized with 1.5% Goat serum/0.1% saponin in PBS for 45' at RT. Incubation of *ER-TR-7* in block&perm solution for 45' at RT. Specific staining was detected with a fluorescent conjugated Goat anti-Rat-IgG (Ref.5).

Positive Control: Spleen.

Negative Control: Lymphoid cells.

Typical Starting Working Dilutions: 1/50 for Immunohistochemistry and Flow Cytometry.

Reactivity: Human, Mouse

Host: Rat

Isotype: IgG2a


Clonality: Monoclonal

Immunogen: Mouse thymic stromal cells.



[View online »](#)

| | |
|-----------------------|---|
| Specificity: | <p>This Monoclonal antibody <i>ER-TR7</i> recognizes with an intracellular component of Mouse Fibroblasts.</p> <p>The <i>ER-TR7</i> antigen is a ubiquitous component of stromal (interstitial) matrix cartilage and of at least some basement membrane zones. The antigen detected is not a basement membrane component, nor any major collagen type or fibronectin. The antigen detected has a wider tissue distribution than reticulin. ER-TR7 detects an intracellular component of fibroblasts. Since ER-TR7 does not react with purified laminin, collagen types I-V, fibronectin, heparin sulfate proteoglycan, entactin or nidogen, it detects a hitherto uncharacterized antigen.</p> <p>The Monoclonal antibody <i>ER-TR7</i> can be used to study the micro-anatomy of various organs. <i>ER-TR7</i> outlines the various compartments of peripheral lymphoid organs by characteristic labeling patterns (no such compartments are found in central lymphoid organs). Furthermore <i>ER-TR7</i> delineates various types of connective tissue compartments in nonlymphoid organs. The antibody <i>ER-TR7</i> detects reticular fibroblasts, which constitute the cellular framework of lymphoid and nonlymphoid organs and their products. <i>ER-TR7</i> is useful to clearly delineated the follicles, periarteriolar lymphoid sheath and marginal zone; the major white pulp compartments. Furthermore in lymph nodes, the capsule, sinuses, follicles, paracortex and medullary cords are clearly delineated.</p> |
| Formulation: | <p>PBS State: Purified State: Liquid 0.2 µm filtered Ig fraction Stabilizer: 0.1% BSA Preservative: 0.02% Sodium Azide</p> |
| Concentration: | lot specific |
| Purification: | Affinity Chromatography on Protein G |
| Conjugation: | Unconjugated |
| Storage: | Store undiluted at 2-8°C. |
| Stability: | Shelf life: one year from despatch. |
| Background: | <p>Fibroblasts are the least specialized cells in the connective-tissue family. They are dispersed in connective tissue throughout the body, where they secrete a nonrigid extracellular matrix (ECM) that is rich in type I and/or type III collagen. Connective tissue consists of glycosaminoglycans, proteoglycans and glycoproteins through which various fibres run. These fibres can be collagenous, elastic or reticular. Reticular fibres are composed from the family of collagen proteins and give tensile strength. These fibres are made by reticular fibroblasts. The activation of fibroblasts by inflammatory stimuli results in their migration, proliferation and deposition of extracellular matrix components, important features involved in both wound healing and fibrosis.</p> |
| Synonyms: | Fibroblast Marker, Fibroblasten |

Product images:

Staining of Mouse C57BL/6 Spleen Section with Antibody Cat.-No BM4018S Clone ER-TR7 at 5 g/ml.