

Product datasheet for BM4047PP

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD31 (PECAM1) Mouse Monoclonal Antibody [Clone ID: MEM-05]

Product data:

Product Type: Primary Antibodies

Clone Name: MEM-05

Applications: FC

Recommended Dilution: Flow Cytometry analysis of human blood cells using 10 μl reagent / 100 μl of whole blood

or 106 cells in a suspension.

The content of a vial (1 ml) is sufficient for 100 tests.

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Leukocytes of patient suffering from LGL-type leukaemia

Specificity: This antibody reacts with CD31 (PECAM-1), a 130-140 kDa type I transmembrane glycoprotein

expressed on monocytes, platelets, granulocytes, endothelial cells and stem cells of the

myeloid lineage.

Formulation: Phosphate buffered saline (PBS)

Label: PerCP

State: Liquid purified lg fraction Preservative: 15 mM sodium azide

Label: Conjugated with Peridinin-chlorophyll-protein complex (PerCP) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for

direct use.

Conjugation: PerCP

Storage: Store undiluted at 2-8°C. DO NOT FREEZE! This products is photosensitive and should be

protected from light.

Stability: Shelf life: one year from despatch.

Gene Name: platelet and endothelial cell adhesion molecule 1

Database Link: Entrez Gene 5175 Human

P16284





CD31 (PECAM1) Mouse Monoclonal Antibody [Clone ID: MEM-05] - BM4047PP

Background:

CD31 (platelet endothelial cell adhesion molecule-1, PECAM-1) is an inhibitory coreceptor involved in regulation of T cell and B cell signaling by a dual immunoreceptor tyrosine-based inhibitory motif (ITIM) that upon associated kinases-mediated phosphorylation provide docking sites for protein-tyrosine phosphatases. CD31 is expressed ubiquitously within the vascular compartment and is located mainly at junctions between adjacent cells. N-terminal Ig-like domain of CD31 is responsible for its homophilic binding, which plays an important role in cell-cell interactions. CD31 is a multifunctional molecule with diverse roles in modulation of integrin-mediated cell adhesion, transendothelial migration, angiogenesis, apoptosis, negative regulation of immunoreceptor signaling, autoimmunity, macrophage phagocytosis, IgE-mediated anaphylaxis and thrombosis. It is one of key regulatory molecules in vascular system.

Synonyms:

PECAM-1, EndoCAM, GPIIA'