

Product datasheet for **BM6012P**

Neurofilament (NEFM) (phosphospecific) Mouse Monoclonal Antibody [Clone ID: RNF406]

Product data:

Product Type:	Primary Antibodies
Clone Name:	RNF406
Applications:	IHC, WB
Recommended Dilution:	Immunoblotting. Flow Cytometry. Immunocytochemistry. Immunohistochemistry on Frozen Sections. Immunohistochemistry on Paraffin-Embedded Tissues. <i>Recommended Dilutions:</i> 1/50-1/100 for Flow Cytometry, and for Immunohistochemistry with avidin-biotinylated horseradish peroxidase complex (ABC) as detection reagent, and 1/100-1/500 for Immunoblotting.
Reactivity:	Guinea Pig, Hamster, Human, Monkey, Rabbit, Xenopus
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Neurofilament preparation of calf brain tissue.
Specificity:	RNF406 reacts exclusively with the phosphorylated isoform of the 160 kD neurofilament protein. Reacts with Human, Rabbit, Hamster, Monkey, Guinea Pig and Xenopus.
Formulation:	PBS with 0.09% Sodium Azide as preservative. State: Purified State: Liquid purified IgG fraction.
Concentration:	lot specific
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freeze-thaw cycles.
Stability:	Shelf life: One year from despatch.
Gene Name:	neurofilament, medium polypeptide



[View online »](#)

Database Link: [Entrez Gene 4741 Human P07197](#)

Background: Like most other intermediate filament proteins (IFPs), the expression of the different neuronal IFPs is both tissue-specific and developmentally regulated. The neurofilament (NF) triplet proteins (70, 160, and 200 kDa) occur in both the central and peripheral nervous system and are normally restricted to neurons. The 70 kDa NF-protein can self-assemble into a filamentous structure, whereas the 160 kDa and 200 kDa NF-proteins require the presence of the 70 kDa NF-protein to co-assemble. All three NF proteins can be detected by immunohistochemical methods at day 9 or 10 after gestation in the mouse embryo. Although IFPs of the neurofilament type are normally restricted to neurons, there are reports on their expression in non-neuronal cells as well. For example, in heart conduction myocytes NF proteins are expressed together with desmin. In tumor pathology ganglioneuroblastomas and some of the other neuroblastomas are strongly positive with the neurofilament antisera. Also, some neuro-endocrine malignancies may show NF positivity. In cell cultures of neural tissues the neurofilament antibodies can monitor in vitro differentiation.

Synonyms: Neurofilament medium polypeptide, NF-M, NEF3, NEFM, Neurofilament 3, (Neuronal Marker)