

Product datasheet for **BM6030P**

Plakophilin 3 (PKP3) Mouse Monoclonal Antibody [Clone ID: 12B11F8]

Product data:

Product Type:	Primary Antibodies
Clone Name:	12B11F8
Applications:	IF, IHC, IP, WB
Recommended Dilution:	Immunocytochemistry (methanol-fixed cells). Immunohistochemistry on Frozen Tissues. Immunoprecipitation. Western blot detection. For optimal results, fixed cells or tissues should be treated with 0.2% Triton-X 100 for 5 min. prior to incubation with primary antibody. <i>Recommended Dilutions:</i> 1/25-1/200 for Immunohistochemistry with avidin-biotinylated horseradish peroxidase complex (ABC) as detection reagent, and 1/100-1/1000 for Immunoblotting applications.
Reactivity:	Canine, Human, Mouse, Porcine
Host:	Mouse
Isotype:	IgG2a
Clonality:	Monoclonal
Immunogen:	Synthetic peptide corresponding to amino acid residues 779-793 of Human Plakophilin-3 coupled to keyhole limpet hemocyanin via an additional cysteine residue at the N-terminus (CKLHRDFAKGYRKED).
Specificity:	This antibody reacts with an epitope located between residues 779-793 in Human Plakophilin-3.
Formulation:	PBS with 0.09% Sodium Azide as preservative. State: Purified State: Liquid purified IgG fraction.
Concentration:	lot specific
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.



[View online »](#)

Gene Name: plakophilin 3

Database Link: [Entrez Gene 11187 Human Q9Y446](#)

Background: Plakophilin-3 is an Armadillo-like protein present in nuclei and desmosomes of epithelial cells [1]. The expression pattern of this protein seems to be largely restricted to epithelial cell types. Plakophilin-3 can be detected along cell borders in a punctuate staining pattern typical for desmosomal proteins. In addition to the desmosomal immunolocalisation, immunostaining was observed as bright nuclear speckles. Thus, like plakophilin-1 and-2, plakophilin-3 displays a dual intracellular localisation in the desmosomal plaque and in the cell nucleus, and therefore is probably involved in signal transduction pathways between the plasma membrane and the nucleus. The human protein has a predicted molecular mass of 87 kD.

Synonyms: Plakophilin-3