

Product datasheet for BM6034P

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Sele (CR2 Region) Rat Monoclonal Antibody [Clone ID: UZ6]

Product data:

Product Type: Primary Antibodies

Clone Name: UZ6

Applications: ELISA, FC, IF, IHC, IP, WB

Recommended Dilution: ELISA.

Immunoblotting. Immunoprecipitation. Flow Cytometry: 1/500. Immunofluorescence. Immunocytochemistry.

Immunohistochemistry on Fozen Sections.

Reactivity: Mouse Host: Rat

Isotype: IgG2b

Clonality: Monoclonal

Immunogen: LPS-activated mlEND1 cells expressing E-selectin at their cell surface.

Specificity: UZ6 recognizes a conformational epitope in the CR2 region in murine E-selectin.

UZ6 partially inhibits leukocyte adhesion to murine activated endothelial cells.

Formulation: PBS with 0.09% Sodium Azide as preservative.

State: Purified

State: Liquid purified IgG fraction.

Concentration: lot specific

Conjugation: Unconjugated

Storage: Store the antibody (undiluted) at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freeze-thaw cycles.

Stability: Shelf life: One year from despatch.

Gene Name: selectin, endothelial cell

Database Link: Entrez Gene 20339 Mouse

Q00690






Background:

Leukocytes adhere to the blood vessel endothelium during extravasation in postcapillary venules of lymph nodes. In addition, leukocyte adhesion occurs in the capillaries and small venules at any site in the body after onset of inflammation. This response is immediate and involves a cascade of adhesion receptors. At the endothelial surface members of the selectin and immunoglobulin superfamilies participate in this cascade. Selectins are C-type cell surface lectins that play a role in leukocyte adhesion to the blood vessel wall endothelium. E-selectin (CD62E) is an endothelial cell specific selectin that is expressed only after activation with proinflammatory cytokines. In vitro experiments have shown that IL-1, TNFalpha and bacterial wall components like lipopolysaccharides induce the transcription of E-selectin in a NFkB dependent signalling cascade. E-selectin has been associated with blood vessel endothelium in diverse inflammatory situations.

Synonyms: SELE, ELAM1, LECAM2

Product images:



Immunostaining of E-Selectin transfected Chinese Hamster Ovary (CHO) cell.