

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **Product datasheet for BP5113**

## Drebrin (DBN1) (324-343) Guinea Pig Polyclonal Antibody

## **Product data:**

Product Type:	Primary Antibodies
Applications:	IF, IHC, IP, WB
Recommended Dilution:	<ul> <li>Immunofluorescence / Immunocytochemistry (assay dependent).</li> <li>Immunoblotting (assay dependent).</li> <li>Immunoprecipitation (assay dependent).</li> <li>Immunohistochemistry on Frozen Paraformaldehyde-Fixed Sections 1/100.</li> <li>Immunohistochemistry on Paraffin Sections: 1/100 (Microwave treatment is recommended)</li> <li>Incubation Time: 1h at RT, extended with paraffin.</li> <li>Immunohistochemistry on Cytological Material.</li> </ul>
Reactivity:	Bovine, Human, Mouse, Rat
Host:	Guinea Pig
Clonality:	Polyclonal
Immunogen:	Synthetic Human peptide (aa 324-343) coupled to KLH.
Specificity:	The antiserum specifically reacts with Drebrin isoforms E2 and A, widespread actin- associating proteins of 70 kD (SDS-PAGE mobility shows Mr 120,000). Reacts with PLC, U333, HUVEC, HaCat, HeLa, MCF-7, SV-80; B1; B16 (Mouse melanoma cells).
Formulation:	State: Serum State: Liquid Serum Stabilizer: 0.5% BSA Preservative: 0.09% Sodium Azide
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	drebrin 1
Database Link:	<u>Entrez Gene 1627 Human</u> <u>Q16643</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	Drebrin (DBN1) (324-343) Guinea Pig Polyclonal Antibody – BP5113
Background:	Drebrin has originally been described only in neuronal cells, but has been recently found also in the actin-rich lamellipodia and filopodia of motile cells. It is enriched at junctional plaques, defining a specific microfilament anchorage system in polar epithelial cells. Positive staining was found in many tissues, including diverse epithelia and carcinoma (e.g. in epidermis basal cells are positive, whereas overlying cells are negative; also positive: regenerating epithelium during wound healing; basal cell carcinoma; skin melanoma), specific types of endothelia and smooth muscle; especially prominent in the mesangial cells of renal glomeruli and in the Sertoli cells of testis. Also positive: epithelia of hair follicles and eccrine sweat glands. Consistently negative were, however, hepatocytes and cross-striated muscle.
Synonyms:	DBN1, DBN-1, D0S117E, DKFZp434D064,drebrin 1

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US