

## **Product datasheet for CF190213**

# OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### **BrdU Mouse Monoclonal Antibody [Clone ID: OTI4G7]**

#### **Product data:**

Isotype:

**Product Type:** Primary Antibodies

lgG1

Clone Name: OTI4G7

Applications: IHC

Recommended Dilution: IHC 1:150

Host: Mouse

Clonality: Monoclonal

Immunogen: Bromodeoxyuridine coupled to keyhole limpet hemocyanin (KLH).

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

**Reconstitution Method:** For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Note: 5-chloro-2'-deoxyuridine (CIDU), 5-bromo-2'-deoxyuridine (BrdU), 5-iodo-2'-deoxyuridine (IdU)


and 5-ethynyl-2'-deoxyuridine (EdU) are nucleoside analogs of thymidine. Cells that treated with these analogs will incorporate the chemicals into the genomic DNA during S-phase. Immunochemical method detection of these analogs is thus used to quantify the cell proliferation, cell cycle status in vitro or in vivo. Since the thymidine analogs can be passed onto the daughter cells, they can also used to trace dividing cell fate in a short period of time over 3 generations. In addition, combination of different analogs and their specific antibodies

can be used to trace cell fate in different time frames.







### **Product images:**



Immunohistochemical staining of paraffinembedded Mouse BRDU colon tissue using anti-BRDU mouse monoclonal antibody. ([TA190213])



Immunohistochemical staining of paraffinembedded colon tissue from IDU injected mouse using anti-BrdU mouse monoclonal antibody. ([TA190213])



Immunohistochemical staining of paraffinembedded colon tissue from CLDU injected mouse using anti-BrdU mouse monoclonal antibody. (TA1905213)