

## Product datasheet for CF190221

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### **BrdU Mouse Monoclonal Antibody [Clone ID: OTI3A8]**

#### **Product data:**

**Product Type: Primary Antibodies** 

IHC

Clone Name: OTI3A8

**Applications:** 

Recommended Dilution: IHC 1:150 Host: Mouse

Isotype: lgG1

Monoclonal Clonality:

Immunogen: Bromodeoxyuridine coupled to keyhole limpet hemocyanin (KLH). Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

**Reconstitution Method:** For reconstitution, we recommend adding 100uL distilled water to a final antibody

> concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Unconjugated Conjugation:

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Note: 5-chloro-2'-deoxyuridine (CIDU), 5-bromo-2'-deoxyuridine (BrdU), 5-iodo-2'-deoxyuridine (IdU)


> and 5-ethynyl-2'-deoxyuridine (EdU) are nucleoside analogs of thymidine. Cells that treated with these analogs will incorporate the chemicals into the genomic DNA during S-phase. Immunochemical method detection of these analogs is thus used to quantify the cell proliferation, cell cycle status in vitro or in vivo. Since the thymidine analogs can be passed onto the daughter cells, they can also used to trace dividing cell fate in a short period of time over 3 generations. In addition, combination of different analogs and their specific antibodies

can be used to trace cell fate in different time frames.







# **Product images:**



Immunohistochemical staining of paraffinembedded colon tissue from BrdU injected mouse using anti-BrdU rat monoclonal antibody ([TA190221], 1:150).



Immunohistochemical staining of paraffinembedded colon tissue from IDU injected mouse using anti-BrdU rat monoclonal antibody ([TA190221], 1:150).



Immunohistochemical staining of paraffinembedded colon tissue from CIDU injected mouse using anti-BrdU rat monoclonal antibody ([TA190221], 1:150).