

Product datasheet for CF500002

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200 Rockville, MD 20850 LIS

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

c-Myc (MYC) Mouse Monoclonal Antibody [Clone ID: OTI1A6]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1A6

Applications: FC, IF, IHC, WB

Recommended Dilution: WB 1:2000, IHC 1:50, IF 1:100, FLOW 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human Myc (NP_002458) produced in E.coli.

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

Reconstitution Method: For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 50.4 kDa

Gene Name: MYC proto-oncogene, bHLH transcription factor

Database Link: NP 002458

Entrez Gene 17869 MouseEntrez Gene 24577 RatEntrez Gene 4609 Human

P01106





Background: c-Myc is a multifunctional, nuclear phosphoprotein that plays a role in cell cycle progression,

apoptosis and cellular transformation. It functions as a transcription factor that regulates transcription of specific target genes. Mutations, overexpression, rearrangement and translocation of this gene have been associated with a variety of hematopoietic tumors, leukemias and lymphomas, including Burkitt lymphoma. The synthesis of non-AUG initiated protein is suppressed in Burkitt's lymphomas, suggesting its importance in the normal

function of this gene.

Synonyms: bHLHe39; c-Myc; MRTL; MYCC

Protein Families: Druggable Genome, Embryonic stem cells, Induced pluripotent stem cells, Stem cell -

Pluripotency, Stem cell relevant signaling - JAK/STAT signaling pathway, Stem cell relevant signaling - TGFb/BMP signaling pathway, Stem cell relevant signaling - Wnt Signaling pathway,

Transcription Factors

Protein Pathways: Acute myeloid leukemia, Bladder cancer, Cell cycle, Chronic myeloid leukemia, Colorectal

cancer, Endometrial cancer, ErbB signaling pathway, Jak-STAT signaling pathway, MAPK signaling pathway, Pathways in cancer, Small cell lung cancer, TGF-beta signaling pathway,

Thyroid cancer, Wnt signaling pathway