

Product datasheet for **CF500367**


USP13 Mouse Monoclonal Antibody [Clone ID: OTI5B9]

Product data:


Product Type:	Primary Antibodies
Clone Name:	OTI5B9
Applications:	IP, WB
Recommended Dilution:	WB 1:2000, IP: 4ug/mL
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full-length protein expressed in 293T cell transfected with human USP13 expression vector
Formulation:	Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)
Reconstitution Method:	For reconstitution, we recommend adding 100uL distilled water to a final antibody concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	97.3 kDa
Gene Name:	ubiquitin specific peptidase 13
Database Link:	NP_003931 Entrez Gene 72607 MouseEntrez Gene 310306 RatEntrez Gene 8975 Human Q92995
Synonyms:	IsoT-3; ISOT3
Protein Families:	Druggable Genome, Protease



[View online »](#)

Product images:


HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY USP13 ([RC202190], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-USP13. Positive lysates [LY401292] (100ug) and [LC401292] (20ug) can be purchased separately from OriGene.



Immunoprecipitation (IP) of USP13 by using TrueMab monoclonal anti-USP13 antibodies (Negative control: IP without adding anti-USP13 antibody.). For each experiment, 500ul of DDK tagged USP13 overexpression lysates (at 1:5 dilution with HEK293T lysate), 2ug of anti-USP13 antibody and 20ul (0.1mg) of goat anti-mouse conjugated magnetic beads were mixed and incubated overnight. After extensive wash to remove any non-specific binding, the immunoprecipitated products were analyzed with rabbit anti-DDK polyclonal antibody.