

Product datasheet for **CF500405**

Chk2 (CHEK2) Mouse Monoclonal Antibody [Clone ID: OTI2F4]

Product data:

| | |
|-------------------------|--|
| Product Type: | Primary Antibodies |
| Clone Name: | OTI2F4 |
| Applications: | FC, IF, WB |
| Recommended Dilution: | WB 1:500~1000, IF 1:50~100, FLOW 1:100 |
| Reactivity: | Human, Dog, Rat, Monkey, Mouse |
| Host: | Mouse |
| Isotype: | IgG2b |
| Clonality: | Monoclonal |
| Immunogen: | Full length human recombinant protein of human CHEK2 (NP_009125) produced in HEK293T cell. |
| Formulation: | Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose) |
| Reconstitution Method: | For reconstitution, we recommend adding 100uL distilled water to a final antibody concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific) |
| Purification: | Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G) |
| Conjugation: | Unconjugated |
| Storage: | Store at -20°C as received. |
| Stability: | Stable for 12 months from date of receipt. |
| Predicted Protein Size: | 60.9 kDa |
| Gene Name: | Homo sapiens checkpoint kinase 2 (CHEK2), transcript variant 1, mRNA. |
| Database Link: | NP_009125 Entrez Gene 50883 Mouse Entrez Gene 114212 Rat Entrez Gene 486338 Dog Entrez Gene 713668 Monkey Entrez Gene 11200 Human O96017 |



[View online »](#)

Background:

In response to DNA damage and replication blocks, cell cycle progression is halted through the control of critical cell cycle regulators. The protein encoded by this gene is a cell cycle checkpoint regulator and putative tumor suppressor. It contains a forkhead-associated protein interaction domain essential for activation in response to DNA damage and is rapidly phosphorylated in response to replication blocks and DNA damage. When activated, the encoded protein is known to inhibit CDC25C phosphatase, preventing entry into mitosis, and has been shown to stabilize the tumor suppressor protein p53, leading to cell cycle arrest in G1. In addition, this protein interacts with and phosphorylates BRCA1, allowing BRCA1 to restore survival after DNA damage. Mutations in this gene have been linked with Li-Fraumeni syndrome, a highly penetrant familial cancer phenotype usually associated with inherited mutations in TP53. Also, mutations in this gene are thought to confer a predisposition to sarcomas, breast cancer, and brain tumors. This nuclear protein is a member of the CDS1 subfamily of serine/threonine protein kinases. Three transcript variants encoding different isoforms have been found for this gene.

Synonyms:


CDS1; CHK2; hCds1; HuCds1; LFS2; PP1425; RAD53

Protein Families:


Druggable Genome, Protein Kinase, Stem cell - Pluripotency

Protein Pathways:


Cell cycle, p53 signaling pathway

Product images:


HEK293T cells transfected with either [RC201278] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-CHEK2 antibody ([TA500405]), and then analyzed by flow cytometry.




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY CHEK2 ([RC201278], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CHEK2. Positive lysates [LY416128] (100ug) and [LC416128] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-CHEK2 monoclonal antibody.




Anti-CHEK2 mouse monoclonal antibody ([TA500405]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY CHEK2 ([RC201278]).




Immunofluorescent staining of HT29 cells using anti-CHEK2 mouse monoclonal antibody ([TA500405]).



Western blot analysis of extracts (10ug) from a mouse cell line and a mouse tissue by using anti-CHEK2 monoclonal antibody (1:200).



Western blot analysis of extracts (10ug) from 9 Human tissue by using anti-CHEK2 monoclonal antibody at 1:200 (1: Testis; 2: Omentum; 3: Uterus; 4: Breast; 5: Brain; 6: Liver; 7: Ovary; 8: Thyroid gland; 9: Colon).



Equivalent amounts of cell lysates (10 ug per lane) of wild-type HeLa cells (WT, Cat# LC810HELA) and CHEK2-Knockout HeLa cells (KO, Cat# [LC810324]) were separated by SDS-PAGE and immunoblotted with anti-CHEK2 monoclonal antibody [TA500405]. Then the blotted membrane was stripped and reprobed with anti-HSP90AB1 antibody ([TA500494]) as a loading control (1:500).