

Product datasheet for CF500482

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

CD4 Mouse Monoclonal Antibody [Clone ID: OTI3B4]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI3B4
Applications: IF, WB

Recommended Dilution: WB 1:2000, IF 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human CD4 (NP_000607) produced in HEK293T

cell.

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

Reconstitution Method: For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 51.1 kDa

Gene Name: CD4 molecule

Database Link: NP 000607

Entrez Gene 920 Human

P01730

Background: Accessory protein for MHC class-II antigen/T-cell receptor interaction. May regulate T-cell

activation. Induces the aggregation of lipid rafts.





Synonyms: CD4mut; IMD79; OKT4D


Protein Families: Adult stem cells, Druggable Genome, ES Cell Differentiation/IPS, Induced pluripotent stem

cells, Transmembrane


Protein Pathways: Antigen processing and presentation, Cell adhesion molecules (CAMs), Hematopoietic cell

lineage, Primary immunodeficiency, T cell receptor signaling pathway

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY CD4 ([RC206453], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CD4. Positive lysates [LY400209] (100ug) and [LC400209] (20ug) can be purchased separately from OriGene.



Anti-CD4 mouse monoclonal antibody ([TA500482]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY CD4 ([RC206453]).