

# **Product datasheet for CF500524**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## SATB1 Mouse Monoclonal Antibody [Clone ID: OTI4E1]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI4E1

**Applications:** IF, IHC, WB

Recommended Dilution: WB 1:2000 IHC 1:50 IF 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full-length protein expressed in 293T cell transfected with human SATB1 expression vector

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

**Reconstitution Method:** For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 86 kDa

Gene Name: SATB homeobox 1

Database Link: NP 002962

Entrez Gene 20230 MouseEntrez Gene 316164 RatEntrez Gene 6304 Human

Q01826

Background: The protein binds to DNA at special AT-rich sequences at nuclear matrix- or scaffold-

associated regions. Thought to recognize the sugar-phosphate structure of double-stranded

DNA.





### SATB1 Mouse Monoclonal Antibody [Clone ID: OTI4E1] - CF500524


Synonyms: DNA-binding protein SATB1; SATB homeobox 1; scaffold-associating DNA's); special AT-rich

sequence binding protein 1; special AT-rich sequence binding protein 1 (binds to nuclear


matrix

**Protein Families:** Druggable Genome, Transcription Factors

### **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY SATB1 ([RC200421], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-SATB1. Positive lysates [LY418976] (100ug) and [LC418976] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffinembedded Adenocarcinoma of breast tissue using anti-SATB1 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.





Immunohistochemical staining of paraffinembedded Adenocarcinoma of ovary tissue using anti-SATB1mouse monoclonal antibody. Heatinduced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Anti-SATB1 mouse monoclonal antibody ([TA500524]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY SATB1 ([RC200421]).