

Product datasheet for **CF501118**

BTK Mouse Monoclonal Antibody [Clone ID: OTI10E10]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI10E10
Applications:	FC, IF, IHC, WB
Recommended Dilution:	WB 1:2000, IHC 1:50, IF 1:100
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG2a
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human BTK (NP_000052) produced in HEK293T cell.
Formulation:	Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)
Reconstitution Method:	For reconstitution, we recommend adding 100uL distilled water to a final antibody concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	76.3 kDa
Gene Name:	Bruton tyrosine kinase
Database Link:	NP_000052 Entrez Gene 12229 Mouse Entrez Gene 367901 Rat Entrez Gene 695 Human Q06187



[View online »](#)


Background: The protein encoded by this gene plays a crucial role in B-cell development. Mutations in this gene cause X-linked agammaglobulinemia type 1, which is an immunodeficiency characterized by the failure to produce mature B lymphocytes, and associated with a failure of Ig heavy chain rearrangement.

Synonyms: AGMX1; AT; ATK; BPK; IMD1; PSCTK1; XLA


Protein Families: Druggable Genome, Protein Kinase

Protein Pathways: B cell receptor signaling pathway, Fc epsilon RI signaling pathway, Primary immunodeficiency


Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY BTK ([RC211582], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-BTK. Positive lysates [LY424947] (100ug) and [LC424947] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffin-embedded Human liver tissue within the normal limits using anti-BTK mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.




Flow cytometric analysis of living 293T cells transfected with BTK overexpression plasmid ([RC211582]), Red)/empty vector ([PS100001], Blue) using anti-BTK antibody ([TA501118]). Cells incubated with a non-specific antibody (Green) were used as isotype control.




Immunofluorescent staining of HT29 cells using anti-BTK mouse monoclonal antibody ([TA501118]).



Anti-BTK mouse monoclonal antibody ([TA501118]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY BTK ([RC211582]).



Western blot analysis of extracts (10ug) from 3 different cell lines by using anti-BTK monoclonal antibody at 1:200 dilution.



Western blot analysis of extracts (10ug) from a mouse cell line and 3 different mouse tissues by using anti-BTK monoclonal antibody (1:200).




Figure from citation: Flow Cytometry analysis of intracellular BTK expression by using anti-BTK antibody (clone OTI10E10) in CD14+ monocytes. Black histograms indicate BTK expression and Gray histograms indicate isotype control. BTK expression in the patient was reduced compared with that in a normal control. [View Citation](#)