

Product datasheet for **CF501174**

BMAL1 (ARNTL) Mouse Monoclonal Antibody [Clone ID: OT1C11]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OT1C11
Applications:	IF, IHC, WB
Recommended Dilution:	WB 1:2000, IHC 1:50, IF 1:100
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human ARNTL(NP_001169) produced in HEK293T cell.
Formulation:	Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)
Reconstitution Method:	For reconstitution, we recommend adding 100uL distilled water to a final antibody concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	68.5 kDa
Gene Name:	Homo sapiens aryl hydrocarbon receptor nuclear translocator like (ARNTL), transcript variant 1, mRNA.
Database Link:	NP_001169 Entrez Gene 11865 Mouse Entrez Gene 29657 Rat Entrez Gene 406 Human
Background:	The protein encoded by this gene is a basic helix-loop-helix protein that forms a heterodimer with CLOCK. This complex binds an E-box upstream of the PER1 gene, activating this gene and possibly other circadian rhythm-associated genes. Three transcript variants encoding two different isoforms have been found for this gene.
Synonyms:	bHLHe5; BMAL1; BMAL1c; JAP3; MOP3; PASD3; TIC




[View online »](#)


Protein Families: Druggable Genome, Transcription Factors

Protein Pathways: Circadian rhythm - mammal


Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY ARNTL (Cat# [RC207870], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ARNTL (Cat# [TA501174]). Positive lysates [LY400474] (100ug) and [LC400474] (20ug) can be purchased separately from OriGene.




Anti-ARNTL mouse monoclonal antibody ([TA501174]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY ARNTL ([RC207870]).




Immunohistochemical staining of paraffin-embedded bladder tissue within the normal limits using anti-ARNTL mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA501174], Dilution 1:50)




Immunohistochemical staining of paraffin-embedded Carcinoma of prostate tissue using anti-ARNTL mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA501174], Dilution 1:50)




Immunohistochemical staining of paraffin-embedded prostate tissue within the normal limits using anti-ARNTL mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA501174], Dilution 1:50)




Immunohistochemical staining of paraffin-embedded pancreas tissue within the normal limits using anti-ARNTL mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA501174], Dilution 1:50)




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of ovary tissue N93ing anti-ARNTL mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA501174], Dilution 1:50)



Immunohistochemical staining of paraffin-embedded lung tissue within the normal limits using anti-ARNTL mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA501174], Dilution 1:50)



Immunohistochemical staining of paraffin-embedded Adenocarcinoma of colon tissue using anti-ARNTL mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA501174], Dilution 1:50)



Immunohistochemical staining of paraffin-embedded colon tissue within the normal limits using anti-ARNTL mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA501174], Dilution 1:50)