

Product datasheet for **CF501241**

H6PD Mouse Monoclonal Antibody [Clone ID: OT1H6]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OT1H6
Applications:	FC, IF, IHC, WB
Recommended Dilution:	WB 1:1000, IHC 1:50, IF 1:100, FLOW 1:100
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human H6PD (NP_004276) produced in HEK293T cell.
Formulation:	Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)
Reconstitution Method:	For reconstitution, we recommend adding 100uL distilled water to a final antibody concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	88.7 kDa
Gene Name:	hexose-6-phosphate dehydrogenase/glucose 1-dehydrogenase
Database Link:	NP_004276 Entrez Gene 9563 Human O95479




[View online »](#)

Background: There are 2 forms of glucose-6-phosphate dehydrogenase. G form is X-linked and H form, encoded by this gene, is autosomally linked. This H form shows activity with other hexose-6-phosphates, especially galactose-6-phosphate, whereas the G form is specific for glucose-6-phosphate. Both forms are present in most tissues, but H form is not found in red cells.


Synonyms: CORTRD1; G6PDH; GDH

Protein Pathways: Metabolic pathways, Pentose phosphate pathway


Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY H6PD ([RC209890], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-H6PD. Positive lysates [LY401369] (100ug) and [LC401369] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffin-embedded Human liver tissue within the normal limits using anti-H6PD mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA501241], Dilution 1:50)




Immunohistochemical staining of paraffin-embedded Carcinoma of Human liver tissue using anti-H6PD mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA501241], Dilution 1:50)




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human ovary tissue using anti-H6PD mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA501241], Dilution 1:50)



Anti-H6PD mouse monoclonal antibody ([TA501241]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY H6PD ([RC209890]).



Immunofluorescent staining of HeLa cells using anti-H6PD mouse monoclonal antibody ([TA501241]).



HEK293T cells transfected with either [RC209890] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-H6PD antibody ([TA501241]), and then analyzed by flow cytometry.