

Product datasheet for CF501354

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

MIOX Mouse Monoclonal Antibody [Clone ID: OTI5H10]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI5H10

Applications: IF, IHC, WB

Recommended Dilution: WB 1:2000, IHC 1:50, IF 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human MIOX(NP_060054) produced in HEK293T

cell

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

Reconstitution Method: For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 32.8 kDa

Gene Name: myo-inositol oxygenase

Database Link: NP 060054

Entrez Gene 55586 Human

Q9UGB7


Synonyms: ALDRL6

Protein Pathways: Ascorbate and aldarate metabolism, Inositol phosphate metabolism







Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY MIOX ([RC210070], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-MIOX. Positive lysates [LY413679] (100ug) and [LC413679] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Human Kidney tissue within the normal limits using anti-MIOX mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA501354], Dilution 1:50)



Anti-MIOX mouse monoclonal antibody ([TA501354]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY MIOX ([RC210070]).