

## Product datasheet for **CF501384**

### **XPNPEP1 Mouse Monoclonal Antibody [Clone ID: OTI1E7]**

#### **Product data:**

|                                |  |
|--------------------------------|--|
| <b>Product Type:</b>           | Primary Antibodies   |
| <b>Clone Name:</b>             | OTI1E7   |
| <b>Applications:</b>           | IF, IHC, WB  |
| <b>Recommended Dilution:</b>   | WB 1:2000, IF 1:100, IHC: 1:150  |
| <b>Reactivity:</b>             | Human, Mouse, Rat  |
| <b>Host:</b>                   | Mouse  |
| <b>Isotype:</b>                | IgG1   |
| <b>Clonality:</b>              | Monoclonal   |
| <b>Immunogen:</b>              | Full length human recombinant protein of human XPNPEP1 (NP_065116) produced in HEK293T cell.   |
| <b>Formulation:</b>            | Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)  |
| <b>Reconstitution Method:</b>  | For reconstitution, we recommend adding 100uL distilled water to a final antibody concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific) |
| <b>Purification:</b>           | Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)  |
| <b>Conjugation:</b>            | Unconjugated   |
| <b>Storage:</b>                | Store at -20°C as received.  |
| <b>Stability:</b>              | Stable for 12 months from date of receipt.   |
| <b>Predicted Protein Size:</b> | 74.6 kDa   |
| <b>Gene Name:</b>              | X-prolyl aminopeptidase 1  |
| <b>Database Link:</b>          | <a href="#">NP_065116</a><br><a href="#">Entrez Gene 170750 Mouse</a> <a href="#">Entrez Gene 170751 Rat</a> <a href="#">Entrez Gene 7511 Human</a><br><a href="#">Q9NQW7</a>  |



[View online »](#)

**Background:**


This gene encodes the cytosolic form of a metalloaminopeptidase that catalyzes the cleavage of the N-terminal amino acid adjacent to a proline residue. The gene product may play a role in degradation and maturation of tachykinins, neuropeptides, and peptide hormones. Alternative splicing results in multiple transcript variants.

**Synonyms:**


APP1; SAMP; XPNPEP; XPNPEPL; XPNPEPL1

**Protein Families:**


Druggable Genome, Protease

**Product images:**


HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY XPNPEP1 ([RC203535], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-XPNPEP1. Positive lysates [LY402779] (100ug) and [LC402779] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffin-embedded Human prostate tissue within the normal limits using anti-XPNPEP1 mouse monoclonal antibody. ([TA501384]). Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Anti-XPNPEP1 mouse monoclonal antibody ([TA501384]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY XPNPEP1 ([RC203535]).