

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for CF501541

UCK (UCK1) Mouse Monoclonal Antibody [Clone ID: OTI7E11]

Product data:

Product Type:	Primary Antibodies		
Clone Name:	OTI7E11		
Applications:	WB		
Recommended Dilution:	WB 1:500		
Reactivity:	Human, Mouse, Rat		
Host:	Mouse		
lsotype:	lgG1		
Clonality:	Monoclonal		
Immunogen:	Full length human recombinant protein of human UCK1 (NP_113620) produced in HEK293T cell.		
Formulation:	Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)		
Reconstitution Method:	For reconstitution, we recommend adding 100uL distilled water to a final antibody concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)		
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)		
Conjugation:	Unconjugated		
Storage:	Store at -20°C as received.		
Stability:	Stable for 12 months from date of receipt.		
Predicted Protein Size:	31.3 kDa		
Gene Name:	uridine-cytidine kinase 1		
Database Link:	<u>NP 113620</u> <u>Entrez Gene 83549 Human</u> <u>Q9HA47</u>		
Background:	Uridine/cytidine kinase-1 (EC 2.7.1.48) is a pyrimidine ribonucleoside kinase that catalyzes the phosphorylation of uridine and cytidine to form uridine monophosphate (UMP) and cytidine monophosphate (CMP) (Van Rompay et al., 2001 [PubMed 11306702]). [supplied by OMIM]		



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

ORIGENE UCK (UCK1) Mouse Monoclonal Antibody [Clone ID: OTI7E11] – CF501541		
Synonyms:	URK1	
Protein Families:	Druggable Genome	
Protein Pathways:	Drug metabolism - other enzymes, Metabolic pathways, Pyrimidine metabolism	
Product images:		

170	_	
130	—	
100	_	
70	_	
55	_	
40	_	
35	—	-
25	—	
15	_	
10	_	

HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY UCK1 ([RC220876], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-UCK1. Positive lysates [LY403114] (100ug) and [LC403114] (20ug) can be purchased separately from OriGene.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US