

Product datasheet for **CF501543**

UCK (UCK1) Mouse Monoclonal Antibody [Clone ID: OTI5E9]


Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI5E9
Applications:	FC, IF, WB
Recommended Dilution:	WB 1:2000, IF 1:100, FLOW 1:100
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human UCK1 (NP_113620) produced in HEK293T cell.
Formulation:	Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)
Reconstitution Method:	For reconstitution, we recommend adding 100uL distilled water to a final antibody concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	31.3 kDa
Gene Name:	uridine-cytidine kinase 1
Database Link:	NP_113620 Entrez Gene 83549 Human Q9HA47
Background:	Uridine/cytidine kinase-1 (EC 2.7.1.48) is a pyrimidine ribonucleoside kinase that catalyzes the phosphorylation of uridine and cytidine to form uridine monophosphate (UMP) and cytidine monophosphate (CMP) (Van Rompay et al., 2001 [PubMed 11306702]). [supplied by OMIM]




[View online »](#)


Synonyms: URK1
Protein Families: Druggable Genome
Protein Pathways: Drug metabolism - other enzymes, Metabolic pathways, Pyrimidine metabolism

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY UCK1 ([RC220876], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-UCK1. Positive lysates [LY403114] (100ug) and [LC403114] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (10ug) from 4 different cell lines by using anti-UCK1 monoclonal antibody (1:200).



Anti-UCK1 mouse monoclonal antibody ([TA501543]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY UCK1 ([RC220876]).



HEK293T cells transfected with either [RC220876] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-UCK1 antibody ([TA501543]), and then analyzed by flow cytometry.