

## Product datasheet for **CF501721**

### PIK3AP1 Mouse Monoclonal Antibody [Clone ID: OTI5H5]

#### Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI5H5
Applications:	FC, IF, WB
Recommended Dilution:	WB 1:2000, IF 1:100, FLOW 1:100
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human PIK3AP1 (NP_689552) produced in HEK293T cell.
Formulation:	Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)
Reconstitution Method:	For reconstitution, we recommend adding 100uL distilled water to a final antibody concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	90.2 kDa
Gene Name:	phosphoinositide-3-kinase adaptor protein 1
Database Link:	<a href="#">NP_689522</a> <a href="#">Entrez Gene 294048 Rat</a> <a href="#">Entrez Gene 118788 Human</a> <a href="#">Q6ZUJ8</a>



[View online »](#)


**Background:** Involved in the activation of phosphoinositide 3-kinase (PI3K) in B-cells and in natural killer (NK) cells. Couples B-cell antigen receptor (BCR) to PI3K activation by providing a docking site for the PI3K subunit PIK3R1, which contributes to B-cell development. Seems to have a complementary role with CD19 in PI3K activation (By similarity). May be involved in the survival of mature B cells via activation of REL (By similarity).

**Synonyms:** BCAP


**Protein Families:** Druggable Genome

**Protein Pathways:** B cell receptor signaling pathway


### Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PIK3AP1 ([RC214125], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PIK3AP1. Positive lysates [LY403461] (100ug) and [LC403461] (20ug) can be purchased separately from OriGene.



Anti-PIK3AP1 mouse monoclonal antibody ([TA501721]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY PIK3AP1 ([RC214125]).



HEK293T cells transfected with either [RC214125] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-PIK3AP1 antibody ([TA501721]), and then analyzed by flow cytometry.