

Product datasheet for **CF501742**

IVD Mouse Monoclonal Antibody [Clone ID: OTI1B10]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI1B10
Applications:	IF, IHC, WB
Recommended Dilution:	WB 1:500, IHC 1:150, IF 1:100, FLOW 1:100
Reactivity:	Human, Monkey, Mouse, Rat, Dog
Host:	Mouse
Isotype:	IgG2b
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human IVD (NP_002612) produced in HEK293T cell.
Formulation:	Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)
Reconstitution Method:	For reconstitution, we recommend adding 100uL distilled water to a final antibody concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	46.5 kDa
Gene Name:	Homo sapiens isovaleryl-CoA dehydrogenase (IVD), transcript variant 1, mRNA; nuclear gene for mitochondrial product.
Database Link:	NP_002216 Entrez Gene 24513 Rat Entrez Gene 56357 Mouse Entrez Gene 100856316 Dog Entrez Gene 702867 Monkey Entrez Gene 3712 Human P26440



[View online »](#)


Background: Isovaleryl-CoA dehydrogenase (IVD) is a mitochondrial matrix enzyme that catalyzes the third step in leucine catabolism. The genetic deficiency of IVD results in an accumulation of isovaleric acid, which is toxic to the central nervous system and leads to isovaleric acidemia. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq]

Synonyms: ACAD2; IVDH


Protein Families: Druggable Genome

Protein Pathways: Metabolic pathways, Valine, leucine and isoleucine degradation


Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY IVD (Cat# [RC201077], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-IVD (Cat# [TA501742]). Positive lysates [LY432239] (100ug) and [LC432239] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-IVD monoclonal antibody.




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human breast tissue using anti-IVD mouse monoclonal antibody ([TA501742]) at 1:150 dilution.




Immunohistochemical staining of paraffin-embedded Human Kidney tissue within the normal limits using anti-IVD mouse monoclonal antibody ([TA501742]) at 1:150 dilution.




Immunohistochemical staining of paraffin-embedded Human liver tissue within the normal limits using anti-IVD mouse monoclonal antibody ([TA501742]) at 1:150 dilution.




Immunohistochemical staining of paraffin-embedded Carcinoma of Human thyroid tissue using anti-IVD mouse monoclonal antibody ([TA501742]) at 1:150 dilution.




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human endometrium tissue using anti-IVD mouse monoclonal antibody ([TA501742]) at 1:150 dilution.




Immunohistochemical staining of paraffin-embedded Carcinoma of Human prostate tissue using anti-IVD mouse monoclonal antibody ([TA501742]) at 1:150 dilution.




Immunohistochemical staining of paraffin-embedded Human bladder tissue within the normal limits using anti-IVD mouse monoclonal antibody ([TA501742]) at 1:150 dilution.



Immunohistochemical staining of paraffin-embedded Carcinoma of Human bladder tissue using anti-IVD mouse monoclonal antibody ([TA501742]) at 1:150 dilution.



Immunohistochemical staining of paraffin-embedded Human lymph node tissue within the normal limits using anti-IVD mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA501742], Dilution 1:50)



Anti-IVD mouse monoclonal antibody ([TA501742]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY IVD ([RC201077]).