

Product datasheet for **CF501841**


MICAL1 Mouse Monoclonal Antibody [Clone ID: OTI1C2]

Product data:


Product Type:	Primary Antibodies
Clone Name:	OTI1C2
Applications:	FC, IHC, WB
Recommended Dilution:	WB 1:2000, FLOW 1:100, IHC 1:150
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human MICAL1 (NP_073602) produced in HEK293T cell.
Formulation:	Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)
Reconstitution Method:	For reconstitution, we recommend adding 100uL distilled water to a final antibody concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	117.7 kDa
Gene Name:	Homo sapiens microtubule associated monooxygenase, calponin and LIM domain containing 1 (MICAL1), transcript variant 1, mRNA.
Database Link:	NP_073602 Entrez Gene 64780 Human Q8TDZ2
Synonyms:	MICAL; MICAL-1; NICAL




[View online >](#)

Product images:


HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY MICAL1 ([RC208308], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-MICAL1. Positive lysates [LY402940] (100ug) and [LC402940] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffin-embedded Human colon tissue within the normal limits using anti-MICAL1 mouse monoclonal antibody ([TA501841]) at 1:150 dilution.




Immunohistochemical staining of paraffin-embedded Human liver tissue within the normal limits using anti-MICAL1 mouse monoclonal antibody ([TA501841]) at 1:150 dilution.




Immunohistochemical staining of paraffin-embedded Human pancreas tissue within the normal limits using anti-MICAL1 mouse monoclonal antibody ([TA501841]) at 1:150 dilution.




Immunohistochemical staining of paraffin-embedded Human thyroid tissue within the normal limits using anti-MICAL1 mouse monoclonal antibody ([TA501841]) at 1:150 dilution.




Immunohistochemical staining of paraffin-embedded Human endometrium tissue within the normal limits using anti-MICAL1 mouse monoclonal antibody ([TA501841]) at 1:150 dilution.




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human endometrium tissue using anti-MICAL1 mouse monoclonal antibody ([TA501841]) at 1:150 dilution.




Immunohistochemical staining of paraffin-embedded Human prostate tissue within the normal limits using anti-MICAL1 mouse monoclonal antibody ([TA501841]) at 1:150 dilution.



Immunohistochemical staining of paraffin-embedded Carcinoma of Human prostate tissue using anti-MICAL1 mouse monoclonal antibody ([TA501841]) at 1:150 dilution.



Immunohistochemical staining of paraffin-embedded Human lymphoma tissue using anti-MICAL1 mouse monoclonal antibody (TA501841) at 1:150 dilution.



Flow cytometric Analysis of HeLa cells, using anti-MICAL1 antibody (TA501841), (Red), compared to a nonspecific negative control antibody (TA50011), (Blue).