

Product datasheet for CF501852

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

ALDH1L1 Mouse Monoclonal Antibody [Clone ID: OTI3B11]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI3B11
Applications: IHC, WB

Recommended Dilution: WB 1:500~2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human ALDH1L1 (NP_036322) produced in

HEK293T cell.

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

Reconstitution Method: For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 98.6 kDa

Gene Name: aldehyde dehydrogenase 1 family member L1

Database Link: NP 036322

Entrez Gene 64392 RatEntrez Gene 10840 Human

<u>075891</u>



ALDH1L1 Mouse Monoclonal Antibody [Clone ID: OTI3B11] - CF501852

Background: The protein encoded by this gene catalyzes the conversion of 10-formyltetrahydrofolate,

NADP, and water to tetrahydrofolate, NADPH, and carbon dioxide. The encoded protein belongs to the aldehyde dehydrogenase family and is responsible for formate oxidation in vivo. Deficiencies in this gene can result in an accumulation of formate and subsequent


methanol poisoning. [provided by RefSeq]


Synonyms: 10-fTHF; 10-FTHFDH; FDH; FTHFD


Protein Families: Druggable Genome

Protein Pathways: One carbon pool by folate

Product images:









Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-ALDH1L1 monoclonal antibody.






Immunohistochemical staining of paraffinembedded Human colon tissue within the normal limits using anti-ALDH1L1 mouse monoclonal antibody. ([TA501852]). Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.




Immunohistochemical staining of paraffinembedded Human Kidney tissue within the normal limits using anti-ALDH1L1 mouse monoclonal antibody. ([TA501852]). Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.




Immunohistochemical staining of paraffinembedded Human liver tissue within the normal limits using anti-ALDH1L1 mouse monoclonal antibody. ([TA501852]). Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.






Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-ALDH1L1 mouse monoclonal antibody. ([TA501852]). Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.




Immunohistochemical staining of paraffinembedded Carcinoma of Human prostate tissue using anti-ALDH1L1 mouse monoclonal antibody. ([TA501852]). Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Immunohistochemical staining of paraffinembedded Carcinoma of Human bladder tissue using anti-ALDH1L1 mouse monoclonal antibody. ([TA501852]). Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.





Immunohistochemical staining of paraffinembedded Human lymph node tissue within the normal limits using anti-ALDH1L1 mouse monoclonal antibody. ([TA501852]). Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.