

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for CF502027

# TTL Mouse Monoclonal Antibody [Clone ID: OTI3G5]

## **Product data:**

Product Type:	Primary Antibodies
Clone Name:	OTI3G5
Applications:	WB
Recommended Dilution:	WB 1:500
Reactivity:	Human, Mouse, Rat
Host:	Mouse
lsotype:	lgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human TTL (NP_714923) produced in HEK293T cell.
Formulation:	Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)
Reconstitution Method:	For reconstitution, we recommend adding 100uL distilled water to a final antibody concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	43.0 kDa
Gene Name:	tubulin tyrosine ligase
Database Link:	<u>NP_714923</u> <u>Entrez Gene 69737 MouseEntrez Gene 171572 RatEntrez Gene 150465 Human</u> <u>Q8NG68</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

### **CRIGENE** TTL Mouse Monoclonal Antibody [Clone ID: OTI3G5] – CF502027

Background:TTL is a cytosolic enzyme involved in the posttranslational modification of alpha-tubulin (see<br/>MIM 602529). Alpha-tubulin within assembled microtubules is detyrosinated over time at the<br/>C terminus. After microtubule disassembly, TTL restores the tyrosine residues and<br/>consequently participates in a cycle of tubulin detyrosination and tyrosination (Erck et al.,<br/>2003 [PubMed 14571137]). [supplied by OMIM]. COMPLETENESS: complete on the 3' end.

#### Synonyms: MGC46235

## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY TTL (Cat# [RC207805], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-TTL(Cat# [TA502027]). Positive lysates [LY403517] (100ug) and [LC403517] (20ug) can be purchased separately from OriGene.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US