

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for CF502054

## CALCOCO2 Mouse Monoclonal Antibody [Clone ID: OTI3F2]

## **Product data:**


Product Type:	Primary Antibodies
Clone Name:	OTI3F2
Applications:	FC, IF, WB
Recommended Dilution:	WB 1:2000, IF 1:100
Reactivity:	Human
Host:	Mouse
lsotype:	lgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human CALCOCO2 (NP_005822) produced in HEK293T cell.
Formulation:	Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)
Reconstitution Method:	For reconstitution, we recommend adding 100uL distilled water to a final antibody concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	52.1 kDa
Gene Name:	calcium binding and coiled-coil domain 2
Database Link:	<u>NP_005822</u> <u>Entrez Gene 10241 Human</u> <u>Q13137</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US CALCOCO2 Mouse Monoclonal Antibody [Clone ID: OTI3F2] - CF502054
Background:
The protein encoded by this gene is a subunit of nuclear domain 10 (ND10) bodies. ND10
bodies are nuclear domains appearing immunohistochemically as ten dots per nucleus. They
are believed to be associated with the nuclear matrix on the basis of their resistance to
nuclease digestion and salt extraction. ND10 proteins are removed from the nucleus by
herpes simplex virus-1 infection and may have a role in viral life cycles. [provided by RefSeq].
COMPLETENESS: complete on the 3' end.

#### Synonyms: NDP52

### **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY CALCOCO2 ([RC203843], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CALCOCO2 (1:500).



Anti-CALCOCO2 mouse monoclonal antibody ([TA502054]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY CALCOCO2 ([RC203843]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Flow cytometric analysis of living 293T cells transfected with CALCOCO2 overexpression plasmid ([RC203843]), Red)/empty vector ([PS100001], Blue) using anti-CALCOCO2 antibody ([TA502054]). Cells incubated with a non-specific antibody (Green) were used as isotype control (1:100).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US