

Product datasheet for CF502055

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CALCOCO2 Mouse Monoclonal Antibody [Clone ID: OTI2B11]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2B11
Applications: FC, IF, WB

Recommended Dilution: WB 1:2000, IF 1:100, FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human CALCOCO2 (NP_005822) produced in

HEK293T cell.

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

Reconstitution Method: For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 52.1 kDa

Gene Name: calcium binding and coiled-coil domain 2

Database Link: NP 005822

Entrez Gene 10241 Human

Q13137






Background:


The protein encoded by this gene is a subunit of nuclear domain 10 (ND10) bodies. ND10 bodies are nuclear domains appearing immunohistochemically as ten dots per nucleus. They are believed to be associated with the nuclear matrix on the basis of their resistance to nuclease digestion and salt extraction. ND10 proteins are removed from the nucleus by herpes simplex virus-1 infection and may have a role in viral life cycles. [provided by RefSeq]. COMPLETENESS: complete on the 3' end.

Synonyms: NDP52

Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY CALCOCO2 ([RC203843], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CALCOCO2. Positive lysates [LY417045] (100ug) and [LC417045] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (10ug) from 5 different cell lines by using anti-CALCOCO2 monoclonal antibody (1:200).





Anti-CALCOCO2 mouse monoclonal antibody ([TA502055]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY CALCOCO2 ([RC203843]).



HEK293T cells transfected with either [RC203843] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-CALCOCO2 antibody ([TA502055]), and then analyzed by flow cytometry.