

Product datasheet for **CF502104**

MYD88 Mouse Monoclonal Antibody [Clone ID: OTI1B4]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI1B4
Applications:	FC, IF, WB
Recommended Dilution:	WB 1:2000, IF 1:100, FLOW 1:100
Reactivity:	Human
Host:	Mouse
Isotype:	IgG2a
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human MYD88 (NP_002459) produced in HEK293T cell.
Formulation:	Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)
Reconstitution Method:	For reconstitution, we recommend adding 100uL distilled water to a final antibody concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	33.1 kDa
Gene Name:	Homo sapiens MYD88 innate immune signal transduction adaptor (MYD88), transcript variant 2, mRNA.
Database Link:	NP_002459 Entrez Gene 4615 Human Q99836



[View online »](#)

Background:

This gene encodes a cytosolic adapter protein that plays a central role in the innate and adaptive immune response. This protein functions as an essential signal transducer in the interleukin-1 and Toll-like receptor signaling pathways. These pathways regulate that activation of numerous proinflammatory genes. The encoded protein consists of an N-terminal death domain and a C-terminal Toll-interleukin1 receptor domain. Patients with defects in this gene have an increased susceptibility to pyogenic bacterial infections. Alternate splicing results in multiple transcript variants. [provided by RefSeq]

Synonyms:


MYD88D

Protein Families:


Druggable Genome

Protein Pathways:


Apoptosis, Toll-like receptor signaling pathway

Product images:


HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY MYD88 ([RC229151], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-MYD88. Positive lysates [LY432175] (100ug) and [LC432175] (20ug) can be purchased separately from OriGene.




Anti-MYD88 mouse monoclonal antibody ([TA502104]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY MYD88 ([RC202253]).



HEK293T cells transfected with either [RC202253] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-MYD88 antibody ([TA502104]), and then analyzed by flow cytometry.



Flow cytometric Analysis of HeLa cells, using anti-MYD88 antibody ([TA502104]), (Red), compared to a nonspecific negative control antibody, (Blue).



Flow cytometric Analysis of Jurkat cells, using anti-MYD88 antibody ([TA502104]), (Red), compared to a nonspecific negative control antibody, (Blue).