

## Product datasheet for **CF502162**


### VSIG2 Mouse Monoclonal Antibody [Clone ID: OTI11B3]

#### Product data:


Product Type:	Primary Antibodies
Clone Name:	OTI11B3
Applications:	FC, IF, WB
Recommended Dilution:	WB 1:500~2000, IF 1:100, FLOW 1:100
Reactivity:	Human
Host:	Mouse
Isotype:	IgG2b
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human VSIG2 (NP_055127) produced in HEK293T cell.
Formulation:	Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)
Reconstitution Method:	For reconstitution, we recommend adding 100uL distilled water to a final antibody concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	34.2 kDa
Gene Name:	V-set and immunoglobulin domain containing 2
Database Link:	<a href="#">NP_055127</a> <a href="#">Entrez Gene 23584 Human</a> <a href="#">Q96IQ7</a>
Synonyms:	2210413P10Rik; CTH; CTXL
Protein Families:	Druggable Genome, Transmembrane




[View online »](#)

**Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY VSIG2 ([RC200170], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-VSIG2. Positive lysates [LY415368] (100ug) and [LC415368] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-VSIG2 monoclonal antibody.



Anti-VSIG2 mouse monoclonal antibody ([TA502162]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY VSIG2 ([RC200170]).



HEK293T cells transfected with either [RC200170] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-VSIG2 antibody ([TA502162]), and then analyzed by flow cytometry.